

# GOOD-WAY VAPONA INSECTICIDE *Dry Bait*

A READY  
PREPARED  
BAIT



## CAUTION

Keep Out of Reach of Children.  
See Right Hand Panel for Complete Precaution  
Statement and Use Directions.

## CAUTION

**HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN.** Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate food, water or other stuffs. Keep out of reach of children. If swallowed induce vomiting immediately and get medical attention. **DO NOT USE IN HOMES. ATROPINE IS AN ANTIDOTE.** Use 2 PAM as a human antidote. This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water.

**DO NOT REUSE CONTAINER. DESTROY BY BURYING WITH WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES.**

## DIRECTIONS FOR USE

**DAIRY BARNs, MILK SHEDs, STABLEs, LIVESTOCK BARNs, LOAFING SHEDs, PIG PENS, AND OUT-DOOR AREAs:** Scatter dry bait at the rate of  $\frac{1}{4}$  lb. per 1,000 sq. ft. on floor or ground areas where flies congregate. Ideal baiting sites are around doorways, feed storage rooms, alleyways, and window sills. On freshly limed floors place treatment on food sacks or on paper. Repeat procedure as necessary for practical fly control.

**CAGED POULTRY AND TURKEYS ON WIRE:** Scatter dry bait at the rate of  $\frac{1}{4}$  lb. per 100 sq. ft. on droppings beneath cages or perches. Repeat at weekly intervals.

**POULTRY HOUSEs:** Scatter dry bait at the rate of  $\frac{1}{4}$  lb. per 100 sq. ft. where flies congregate but keep out of reach of birds. Apply to manure on window sills, outside of penned enclosures and on floor of feed storage room. For caged units apply to dropping area. Repeat as necessary.

**DOG KENNELs:** Scatter dry bait at the rate of  $\frac{1}{4}$  lb. per 1,000 sq. ft. outside dog runways, on window sills, ledges and other surfaces out of reach of animals. Repeat as necessary.

**GARBAGE CANS:** Apply bait around and inside garbage can. Repeat as necessary.

**FOR REDUCING POPULATION OF RESISTANT AND NON RESISTANT HOUSEFLIES IN AND AROUND OUTDOOR LIVING AREAs, PICNIC GROUNDS, BACKYARD AREAs, PATIOS, LOADING DOCKs, AND OUTDOOR LATRINES, PARKING AREAs, REFUSE AREAs, AROUND SERVICE STATIONS, OPEN AIR DRIVE-INS, ICE CREAM STANDs, GARBAGE COLLECTION AND DISPOSAL AREAs:** Use dry bait at  $\frac{1}{2}$  lb. per 1,000 sq. ft. of area. Apply on ground or areas where flies congregate. For open air drive-ins, ice cream stands and restaurants, apply on apron between serving counter and parking area, paying particular attention to trash barrels and garbage cans where flies congregate.

ACCEPTED  
JUL 13 1954

OFFICIAL STATEMENT

# GOOD-WAY

*Dry Bait*

**NEW  
WITH VAPONA®  
INSECTICIDE (DDVP)  
USE WET OR DRY  
FAST FLY CONTROL**

ACTIVE INGREDIENTS: By Weight  
2,2-Dichlorovinyl  
dimethyl phosphate 0.485%  
Related Compounds 0.835%  
INERT INGREDIENTS 98.680%

Total 100.000%

E.P.A. Reg. No. 2006-39  
E.P.A. Est. No. 2006-IL-1

\*Equivalent to 0.5% of Vapona  
Insecticide

® Trademark Shell Chemical Co.

**NET CONTENTS 1 LB.**

Distributed by  
**GOOD-WAY  
INSECTICIDE, INC.**  
Wheeling, Illinois

