

# Pellets GOOD-WAY Rat & Mouse Killer

Anticoagulant rodenticide



Kills Norway Rats, Roof Rats and House Mice.

**ACTIVE INGREDIENTS:**

Warfarin, (3(a-acetonylbenzyl)-4-

hydroxycoumarin)

**INERT INGREDIENTS**

0.025%

99.975%

TOTAL 100.000%

KEEP OUT OF REACH OF CHILDREN

## CAUTION:

SEE BACK PANEL FOR COMPLETE CAUTION  
STATEMENT AND ADDITIONAL INFORMATION

### PRECAUTIONARY STATEMENTS: HAZARD TO HUMANS & DOMESTIC ANIMALS

**CAUTION:** Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

**NOTE TO PHYSICIAN** — If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally, as indicated in bis-hydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

### ENVIRONMENTAL HAZARD:

Keep Out of Lakes, Streams or Ponds.

MANUFACTURED BY  
GOOD-WAY INSECTICIDE, INC.  
WHEELING, ILLINOIS

NET WT. 1 LB. (454g.)

## Back Panel

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**USE RESTRICTIONS** — For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

**SELECTION OF TREATMENT AREAS** — Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

### APPLICATION DIRECTIONS

**RATS:** Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

**MICE:** Apply ¼ to ½ ounce (1-2 level table-spoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.


**RATS & MICE:** Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

### STORAGE AND DISPOSAL

**Storage:** Store only in original container, in a dry place inaccessible to children and pets.

**Disposal:** Do not reuse empty container. Securely wrap in newspaper and discard in trash.

EPA Reg. No. 2006-4-3  
EPA Est. No. 2006-IL-1



**BEST AVAILABLE COPY**

# Pellets

## GOOD-WAY

### Rat & Mouse Killer

Anticoagulant rodenticide



**NOT REVIEWED**

In Accordance with EPA Notice 82-2,

Based on labeling changed 1-21-86

ACTIVE INGREDIENTS

Warfarin, 3 active ingredients

40% active ingredient

INERT INGREDIENTS

0.0000%

99.9750%

TOTAL 100.0000%

**KEEP OUT OF REACH OF CHILDREN**

### CAUTION:

SEE BACK PANEL FOR COMPLETE CAUTION STATEMENT AND ADDITIONAL INFORMATION

PRECAUTIONARY STATEMENTS:

HAZARD TO HUMANS & DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

**ENVIRONMENTAL HAZARD:**

Keep Out of Lakes, Streams or Ponds.

MANUFACTURED BY

**GOOD-WAY INSECTICIDE, INC.**

WHEELING, ILLINOIS

**NET WT. 1 LB. (454 G.)**

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**USE RESTRICTIONS-** For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

*Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.*

**SELECTION OF TREATMENT AREAS -** Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in cracks and concealed places, between floors and walls, in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove mice or feed as possible.

### APPLICATION DIRECTIONS

**RATS:** Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

**MICE:** Apply 1/4 to 1 ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 10 days.

**RATS & MICE:** Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

### STORAGE AND DISPOSAL

**STORAGE:** Store only in original container, in a dry place inaccessible to children and pets.

**DISPOSAL:** Do not reuse empty container. Securely wrap in newspaper and discard in trash.

EPA Reg. NO. 2006 - 43

EPA Est. NO. 2006 -IL- 1

**BEST AVAILABLE COPY**