

GOOD-WAY

TOMATO & POTATO Dust or Spray

KILLS
HORN WORMS,
COLORADO POTATO
BEETLES, FLEA
BEETLES, TOMATO
FRUIT WORMS,
LEAFHOPPERS,
TARNISHED PLANT
BUG, EARLY BLIGHT
and LATE BLIGHT.



CAUTION

Keep Out of
Reach of Children

See Back Panel for
Complete Precautionary
Statement and Use

ACTIVE INGREDIENTS:	BY WT.
Carbaryl (1-Naphthyl N-Methylcarbamate)	5%
Copper as Metallic (From Basic Copper Sulphate)	7%
INERT INGREDIENTS:	88%
TOTAL	100%

Contains SEVIN® Carbaryl Insecticide
*Registered trade mark of Union Carbide Corporation for the
active ingredient 1-Naphthyl N-Methylcarbamate

NET CONTENTS: ONE POUND

GOOD-WAY Tomato and Potato Dust or Spray is an insecticidal and fungicidal dust for use against certain insects and plant diseases on Tomatoes and Potatoes.
READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

GOOD-WAY Tomato and Potato Dust or Spray

CAUTION

Harmful if swallowed or absorbed through the skin. Skin contact may be harmful. Avoid breathing dust or spray mist. Wash thoroughly after use.

Avoid contamination of food and foodstuffs. Do not apply to food crops unless residues are washed or removed by other effective means. Do not use on forage crops to be fed to dairy animals or to animals being finished for slaughter. Do not use on cucurbits.

DO NOT DUST OPEN BLOSSOMS. TOXIC TO HONEY BEES.

NOTE TO PHYSICIANS: Carbaryl is a moderate, reversible, Cholinesterase Inhibitor. Atropine is antidotal.

DIRECTIONS

For Disease Control Begin Application
Before Disease Appears

AS A DUST: On young plants, apply at rate of 1/2 lb. to 1,000 sq. ft. 20 lbs. per acre.

AS A SPRAY: Mix 6 to 8 tablespoonfuls in a little water to make a thin paste, add to 1 gallon water and agitate during application.

TOMATOES: Horn Worms, Colorado Potato Beetles, Flea Beetles and Tomato Fruit Worm, Early and Late Blight application every week or 10 days is recommended.

Commence when plants leaf out and continue above during early growing season. Late season application should be applied when necessary. Do not apply within 1 day of harvest.

POTATOES: Leafhoppers, Colorado Potato Beetles, Flea Beetle and Tarnished Plant Bug, Early Blight and Late Blight application every week or 10 days is recommended.

Commence when plants leaf out and continue above during early growing season. Late season application should be applied when necessary. Do not apply within 1 day of harvest.

MANUFACTURED FOR

GOOD-WAY INSECTICIDE, INC.

Wheeling, Illinois

U.S.A. Reg. No. 2006-57

ACCEPTED

JUN 30 1969