



JUL 9 1975
 Under Co. Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under **2006-59**

FOR THE CONTROL OF CERTAIN PESTS ON VEGETABLES, SMALL FRUITS, ORNAMENTALS AND PETS

Active Ingredients:
 Carbaryl (1-naphthyl N-methylcarbamate).....5% by wt.
Inert Ingredients:.....95% by wt.

READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.

*Registered trademark of Union Carbide Corporation for the active ingredient 1-naphthyl N-methylcarbamate.

CAUTION! KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed, inhaled or absorbed through the skin. Avoid Skin Contact. Wash Hands Before Eating. Note for Physician: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal. Avoid contamination of food. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NET WEIGHT 1 LB.

GENERAL DIRECTIONS FOR USE

Apply Good-Way 5% Sevin Dust when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surface of all foliage for complete coverage. Repeat as necessary, generally at 7-day intervals. For best results, apply with a dust gun. Do not apply in excess.

SMALL FRUIT INSECT CONTROL

GRAPES: To control grape berry moth, Japanese beetles, June beetles, red-banded leaf roller, grape leaf folder, grape leafhopper and grape leaf skeletonizer. Apply just before first brood leaf folder larvae emerge from rolls and as needed for leafhoppers. (Treat upper and lower surface of all foliage for complete coverage.) Limitation day of harvest. Tolerance 10 ppm.
STRAWBERRIES: To control meadow spittlebug, strawberry leaf roller and strawberry weevil. (Treat upper and lower surface of all foliage for complete coverage.) Limitation 1 day before harvest. Tolerance 10 ppm.
BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES and RASPBERRIES: To control Japanese beetles, leaf rollers, leafhoppers and raspberry aphid. Treat upper and lower surface of all foliage for complete coverage. Limitations 7 days before harvest. Tolerance 12 ppm.

PET USE

ON DOGS AND CATS: For ticks and fleas dust liberally over the entire animal, avoiding the eyes, and rub thoroughly into the skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Avoid treatment of kittens and puppies less than 4 weeks of age.
IN AND AROUND DOG AND CAT QUARTERS: For ticks and fleas dust liberally in and around sleeping quarters, kennels, and dog houses by means of a shaker can, hand or dust gun.

CROP	INSECTS CONTROLLED	DOSEAGE	LIMITATIONS
Asparagus	On Spears: Asparagus beetle On Ferns And Brush: Asparagus beetle, Apache cicada	Treat upper and lower surface of all foliage for complete coverage.	1 day before harvest. Tolerance 10 ppm.
Olives	Corn earworm, stink bugs.		Day of harvest, Tolerance 10 ppm.
Peas*	Leafhoppers, Colorado potato beetle, Armyworms, grasshoppers.		Day of harvest, Tolerance 10 ppm on peas, 100 ppm on fangs.
Cauliflower*, Broccoli*, Kohlrabi*, Brussels sprouts*, Cabbage*	Flea beetle, harlequin bug, Corn earworms, armyworms, cabbage caterpillars		3 days before harvest. Tolerance 10 ppm.
Potato Eggplant Tomato Pepper	Colorado potato beetle, flea beetles, leaf hoppers Tomato fruitworm, fall armyworm, European corn borer, stink bugs, lace bugs and tarnished plant bugs		Day of harvest, Tolerance 10 ppm. (No tolerance on potatoes.)
Cucurbits*, Casaba Cucumber, Crenshaw, Cantaloupe, Honey Bell, Pumpkin Squash, Muskmelon, Honeydew, Watermelon.	Pickleworm, melonworm Cucumber beetles, leafhoppers, flea beetles, squash bugs		Day of harvest, Tolerance 10 ppm. (Do not use on watermelon in Florida.)
Beans*	Mexican bean beetle Leafhoppers, bean leaf beetle, Western bean cut worm, flea beetles, Japanese beetle, velvet bean caterpillar Corn earworm, armyworms, stink bugs, tarnished plant bug In California: Lima bean pod borer, corn earworm, Lygus and stink bugs		Day of harvest, Tolerance 10 ppm on beans, 100 ppm on forage or hay.
Beet Crops*, Beets Rutabaga Do not apply 14 days before harvest on beets, salsify or turnips if tops are to be used for food purposes.	Flea beetles, leafhoppers, harlequin bug Six-spotted leafhopper (aster yellows vector) Corn earworm, armyworms, cabbage caterpillars, tarnished plant bugs and stink bugs		Day of harvest on carrots, 3 days before harvest on other root crops. Tolerance 10 ppm for carrots, 5 ppm for other root crops.
Leafy Vegetables*, Kale Mustard Greens, Turnip Greens Dandelion, Endive, Swiss Chard, Beet Greens, Spinach, Parsley, Lettuce, Collards, Chinese Cabbage	Flea beetles, leafhoppers, harlequin bug Six-spotted leafhopper (Aster yellows vector) Corn earworm, armyworms, cabbage, caterpillars, tarnished plant bugs and stink bugs		3 days before harvest of head lettuce, 14 days before harvest of other leafy crops. Tolerance 10 ppm for head lettuce, salsify tops, endive, Chinese cabbage; 12 ppm for other leafy greens.
Corn*	Corn earworm, corn rootworm adults, armyworm, sap beetles, Japanese beetle, flea beetles, leafhoppers		Day of harvest, Tolerance 5 ppm in ears, Tolerance 100 ppm on forage and fodder.

For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 or 3-day intervals starting when first silks appear, and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are essential for effective control.

*When label directions for use are followed, forage, vines, hay and crop residues may be fed to meat or dairy animals.

ORNAMENTAL INSECT CONTROL: Good-Way 5% Sevin Dust can be safely used on a wide range of ornamental plants, flowers, shrubs and small trees, such as rose, carnation, gladiolus, chrysanthemum, lilac, arbor vitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. Good-Way 5% Sevin Dust injures Boston ivy. Apply to the upper and lower leaf surface of all foliage for complete coverage to control such pests as:

apple aphid mealy bugs Japanese beetle boxwood leaf miner scale insects plant bugs willow leaf beetles rose-slug
 bagworm mimosa webworm June beetles elm leaf aphid tent caterpillars puss caterpillars periodical cicada leaf roller
 birch leaf miner oak leaf miners lace bugs elm leaf beetle thorn bugs psyllids boxelder bug
 blister beetles orange tortrix leafhoppers flea beetles thrips (exposed) rose aphid gypsy moth

NOTE: Good-Way 5% Sevin Dust does not control spider mites. Where infestations are encountered, use a recommended miticide. For protection of honeybees avoid use, if possible, during periods when honeybees are visiting the crop. When necessary to use during such periods, warn beekeepers will in advance to locate hives at a safe distance until one week after application.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Extension Service. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

Injury to tender foliage may occur if applications are made when plants are wet or in the presence of high humidity of several days duration following treatment. AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS, POND, FEEDING TROUGHS AND WATERING RECEPTACLES.

Distributed by Good-Way Insecticide Inc., Wheeling, Illinois 60090

EPA REG. NO. 2006-59 MADE IN U.S.A.
 E. P. A. EST. NO. 419-1L-1

GOOD-WAY

5% SEVIN DUST

INSECTICIDE
ACCEPTED
Contains SEVIN* Carbaryl Insecticide

JUL 9 1975
United States Federal Insecticide, Fungicide and Disinfectant Act.
This product is a registered pesticide
under the Act.

FOR THE CONTROL OF CERTAIN INSECTS ON VEGETABLES, SMALL FRUITS, ORNAMENTALS AND PETS

Active Ingredients:

Carbaryl (1-naphthyl N-methylcarbamate) 5% by wt.

Inert Ingredients: 95% by wt.

READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.

*Registered trademark of Union Carbide Corporation for the active ingredient 1-naphthyl N-methylcarbamate.

CAUTION! KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed, inhaled or absorbed through the skin. Avoid Skin Contact. Wash Hands Before Eating. Note for Physician: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal. Avoid contamination of food. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NET WEIGHT 1 LB

GENERAL DIRECTIONS FOR USE

Apply Good Way 5% Carbaryl Insecticide Dust to the insects, plants, or animals to be treated. For use on insects, apply dust to the insects. For use on plants, apply dust to the plants. For use on animals, apply dust to the animals. For best results, apply with a dust gun. Do not breathe dust.

SMALL FRUIT INSECT CONTROL

GRAPES - Apply to clusters of grapes, leaves, and vines. For control of grape leaf miner, grape leaf hopper and grape leaf skeletonizer, apply to all surfaces of leaves and vines. For control of grape leaf miner, apply to the upper and lower surface of all foliage for complete coverage.

STRAWBERRIES - Apply to leaves, fruit, and runners. For control of strawberry weevil, apply to the upper and lower surface of all foliage for complete coverage.

BLACKBERRIES, RASPBERRIES, BLUEBERRIES, AND RASPBERRIES - To control Japanese beetles, leaf miners, leafhoppers and raspberry aphids. For control of Japanese beetles, apply to the upper and lower surface of all foliage for complete coverage. Limitations: 7 days before harvest. Tolerance: 12 ppm.

PET USE

ON DOGS AND CATS - For flea and tick control, apply dust to the back of the neck, between the eyes, and rub thoroughly into the skin. Comb out dead tucks a few hours after treatment. For control of fleas and ticks, apply dust to the back of the neck, between the eyes, and rub thoroughly into the skin. For control of fleas and ticks, apply dust to the back of the neck, between the eyes, and rub thoroughly into the skin. For control of fleas and ticks, apply dust to the back of the neck, between the eyes, and rub thoroughly into the skin. For control of fleas and ticks, apply dust to the back of the neck, between the eyes, and rub thoroughly into the skin.

CROP	INSECTS CONTROLLED	DOSAGE	LIMITATIONS
Asparagus	On Spears: Asparagus beetle On Ferns And Brush: Asparagus beetle, Awned leafhopper		1 day before harvest. Tolerance 10 ppm.
Okra	Corn earworm, stink bugs		Day of harvest, Tolerance 10 ppm.
Peas*	Leafhoppers, Curculionids, potato beetle, Armyworms, grasshoppers		Day of harvest, Tolerance 10 ppm on peas, 100 ppm on forage
Cauliflower*, Broccoli*, Kohlrabi*	Flea beetles, leafhoppers, budworms, armyworms, cabbage rootworms		3 days before harvest. Tolerance 10 ppm.
Brussels sprouts*, Cabbage*	Armyworms, cabbage rootworms		Day of harvest, Tolerance 10 ppm. (No tolerance on potatoes.)
Potato Eggplant	Colorado potato beetle, leaf beetles and fleas		Day of harvest, Tolerance 10 ppm. (No tolerance on potatoes.)
Tomato	Tomato fruitworm, fall armyworm, European cornborer, stink bugs, lace bugs and tin snipe, plant bugs		Day of harvest, Tolerance 10 ppm. (Do not use on watermelon in Florida.)
Pepper	Stink bugs, lace bugs and tin snipe, plant bugs		Day of harvest, Tolerance 10 ppm. (Do not use on watermelon in Florida.)
Cucurbits* Cucumber, Melon, Pumpkin, Squash	Pumpkin beetle, leafhoppers, flea beetles, squash bug		Day of harvest, Tolerance 10 ppm. (Do not use on watermelon in Florida.)
Cantaloupe, Honeydew, Watermelon	Flea beetles, squash bug		Day of harvest, Tolerance 10 ppm. (Do not use on watermelon in Florida.)
Beans*	Mexican bean beetle, Leafhoppers, bean leaf beetle, western bean cutworm, flea beetles, Japanese beetle, velvet bean caterpillar, Corn earworm, armyworms, stink bugs, shield plant bug In California: Lima bean pod borer, cutworms, Lygus and stink bugs		Day of harvest, Tolerance 10 ppm on beans, 100 ppm on forage or hay.
Beet Crops*, Beets	Flea beetles, leafhoppers, ground squirrels, flea beetles, Japanese beetle, velvet bean caterpillar		Day of harvest on carrots, 3 days before harvest on other root crops. Tolerance 10 ppm for carrots, 5 ppm for other root crops.
Radish, Carrot	Six spotted leafhopper, velvet bean caterpillar		Day of harvest on carrots, 3 days before harvest on other root crops. Tolerance 10 ppm for carrots, 5 ppm for other root crops.
Salsfy, Turnips	Corn earworm, armyworms, shield plant bug		Day of harvest on carrots, 3 days before harvest on other root crops. Tolerance 10 ppm for carrots, 5 ppm for other root crops.
Parsnips, Horseradish	Shield plant bug and stink bugs		Day of harvest on carrots, 3 days before harvest on other root crops. Tolerance 10 ppm for carrots, 5 ppm for other root crops.
Rutabaga	Shield plant bug and stink bugs		Day of harvest on carrots, 3 days before harvest on other root crops. Tolerance 10 ppm for carrots, 5 ppm for other root crops.
Do not apply 14 days before harvest on beets, salsfy, turnips, or rutabaga to be used for food purposes.			
Leafy Vegetables*, Kale	Flea beetles, leafhoppers, ground squirrels, flea beetles, Japanese beetle, velvet bean caterpillar		3 days before harvest of head lettuce, 14 days before harvest of other leafy crops. Tolerance 10 ppm for head lettuce, salsfy tops, endive, Chinese cabbage, 12 ppm for other leafy greens.
Mustard Greens, Turnip Greens	Six spotted leafhopper, velvet bean caterpillar		3 days before harvest of head lettuce, 14 days before harvest of other leafy crops. Tolerance 10 ppm for head lettuce, salsfy tops, endive, Chinese cabbage, 12 ppm for other leafy greens.
Dandelion, Endive, Swiss Chard	Corn earworm, armyworms, shield plant bug		3 days before harvest of head lettuce, 14 days before harvest of other leafy crops. Tolerance 10 ppm for head lettuce, salsfy tops, endive, Chinese cabbage, 12 ppm for other leafy greens.
Beet Greens, Spinach	Corn earworm, armyworms, shield plant bug		3 days before harvest of head lettuce, 14 days before harvest of other leafy crops. Tolerance 10 ppm for head lettuce, salsfy tops, endive, Chinese cabbage, 12 ppm for other leafy greens.
Parsley, Lettuce, Collards	Stink bugs		3 days before harvest of head lettuce, 14 days before harvest of other leafy crops. Tolerance 10 ppm for head lettuce, salsfy tops, endive, Chinese cabbage, 12 ppm for other leafy greens.
Chinese Cabbage	Stink bugs		3 days before harvest of head lettuce, 14 days before harvest of other leafy crops. Tolerance 10 ppm for head lettuce, salsfy tops, endive, Chinese cabbage, 12 ppm for other leafy greens.
Corn*	Corn earworm, fall armyworm, European cornborer, stink bugs, leafhoppers		Day of harvest, Tolerance 5 ppm in ears, Tolerance 100 ppm on forage and fodder.

For larvae in whorl (budworm) damage, apply to the plants, or animals to be treated. For insects attacking birds and bees, apply at 2 or 3 day intervals during the season. For insects attacking mammals, apply at 2 or 3 day intervals during the season. For insects attacking birds and bees, apply at 2 or 3 day intervals during the season. For insects attacking mammals, apply at 2 or 3 day intervals during the season. For insects attacking birds and bees, apply at 2 or 3 day intervals during the season. For insects attacking mammals, apply at 2 or 3 day intervals during the season.

*When label directions for use are followed, this product is safe for use on dairy animals.

ORNAMENTAL INSECT CONTROL - Good Way 5% Carbaryl Insecticide Dust is effective against a wide range of ornamental plants, flower shrubs and small trees, such as rose, carnation, geranium, petunia, verbena, zinnia, gladiolus, iris, hydrangea, lily, and pine. Good Way 5% Carbaryl Insecticide Dust is effective against a wide range of ornamental plants, flower shrubs and small trees, such as rose, carnation, geranium, petunia, verbena, zinnia, gladiolus, iris, hydrangea, lily, and pine. Good Way 5% Carbaryl Insecticide Dust is effective against a wide range of ornamental plants, flower shrubs and small trees, such as rose, carnation, geranium, petunia, verbena, zinnia, gladiolus, iris, hydrangea, lily, and pine.

NOTE - Good Way 5% Carbaryl Insecticide Dust is effective against a wide range of ornamental plants, flower shrubs and small trees, such as rose, carnation, geranium, petunia, verbena, zinnia, gladiolus, iris, hydrangea, lily, and pine. Good Way 5% Carbaryl Insecticide Dust is effective against a wide range of ornamental plants, flower shrubs and small trees, such as rose, carnation, geranium, petunia, verbena, zinnia, gladiolus, iris, hydrangea, lily, and pine.

This product is highly flammable. Keep away from fire and heat. For information on fire hazards, contact your local fire department. For information on fire hazards, contact your local fire department. For information on fire hazards, contact your local fire department.

Injury to birds, fish, and aquatic life may occur if the dust is applied to water. Do not apply to water. Do not apply to water. Do not apply to water. Do not apply to water. Do not apply to water.

WATER - ACCEPTABLE. WATER - ACCEPTABLE. WATER - ACCEPTABLE. WATER - ACCEPTABLE. WATER - ACCEPTABLE.

**AY
UST**

**PTED
Insecticide**

1975
Insecticide
Rodenticide
pesticide
**ON VEGETABLES
ID PETS**

5% by wt.
95% by wt.

**ACCORDANCE
ONS.**

**Corporation for
arbamate.**

OF CHILDREN.

through the skin.
ng. Note for Phy-
Cholinesterase in-
mination of food.
urrying with waste

GENERAL DIRECTIONS FOR USE

Apply Good Way 5:50 (Dust) with insecticide in the evening or early in the morning, or evening when it is cool. Treat upper and lower surfaces of foliage for complete coverage. Repeat as necessary, generally at 7 day intervals. For best results, apply with a dust gun. Do not apply to birds.

SMALL FRUIT INSECT CONTROL

GRAPES - To control Japanese beetles, leafhoppers, leafminer, grape leafhopper and grape leaf skeletonizer. Apply 1-2 days before first harvest. Repeat as necessary, as often as needed for leafhoppers. Treat upper and lower surface of all foliage for complete coverage. Limitations: 7 days before harvest. Tolerance 10 ppm.

STRAWBERRIES - To control meadow spittlebugs, strawberry rootworm, strawberry weevil. Treat upper and lower surface of all foliage for complete coverage. Limitations: 7 days before harvest. Tolerance 10 ppm.

BLACKBERRIES, BOYSENBERRIES, RASPBERRIES, LOGANBERRIES - To control Japanese beetles, leafminers, leafhoppers and raspberry aphids. Treat upper and lower surface of all foliage for complete coverage. Limitations: 7 days before harvest. Tolerance 12 ppm.

PET USE

ON DOGS AND CATS - For ticks and fleas, rub 1/2 tsp. of the concentrate on the face, neck, ears, and rub thoroughly into the skin. Comb out dead ticks a few hours after treatment. Repeat as necessary. Avoid treatment of kittens and puppies less than 4 weeks of age.
IN AND AROUND DOG AND CAT QUARTERS - For ticks and flea dust, liberally in and around sleeping quarters, kennels, and dog houses by means of a shaker can, hand or dust gun.

CROP	INSECTS CONTROLLED	DOSEAGE	LIMITATIONS
Asparagus	On Spears: Asparagus beetle On Ferns And Brush: Asparagus beetle, Apache cicada	Treat upper and lower surface of all foliage for complete coverage.	1 day before harvest. Tolerance 10 ppm.
Okra	Corn earworm, stink bugs.		Day of harvest, Tolerance 10 ppm.
Peas*	Leafhoppers, Colorado potato beetle, Armyworms, grasshoppers		Day of harvest, Tolerance 10 ppm on peas, 100 ppm on forage.
Cauliflower*, Broccoli*, Kale*, Brussels sprouts*, Cabbage*	Flea beetle, harlequin bug, Corn earworms, armyworms, cabbage caterpillars		3 days before harvest. Tolerance 10 ppm.
Potato Eggplant	Colorado potato beetle, flea beetles, leaf hoppers		Day of harvest, Tolerance 10 ppm. (No tolerance on potatoes.)
Tomato Pepper	Tomato fruitworm, fall armyworm, European corn borer, stink bugs, lace bugs and tarnished plant bugs		Day of harvest, Tolerance 10 ppm. (Do not use on watermelon in Florida.)
Cucurbits* - Cucumber, Crenshaw, Cantaloupe, Honey Ball, Pumpkin Squash, Muskmelon, Honeydew, Watermelon.	Pink worm, melonworm, Cucumber beetles, leafhoppers, flea beetles, squash bugs		Day of harvest, Tolerance 10 ppm on beans, 100 ppm on forage or hay.
Beans*	Mexican bean beetle, Leafhoppers, bean leaf beetle, Western bean cut worm, flea beetles, Japanese beetle, velvet bean caterpillar, Corn earworm, armyworms, stink bugs, tarnished plant bug In California: Lima bean pod borer, corn earworm, Lygus and stink bugs		Day of harvest on carrots, 3 days before harvest on other root crops. Tolerance 10 ppm for carrots, 5 ppm for other root crops.
Beet Crops*, Beets Radish, Carrots, Salsify, Turnips, Parsnips, Horseradish, Rutabaga	Flea beetles, leafhoppers, harlequin bug, Six spotted leafhopper (aster yellows vector), Corn earworm, armyworms, cabbage caterpillar, tarnished plant bugs and stink bugs.		3 days before harvest of head lettuce, 14 days before harvest of other leafy crops. Tolerance 10 ppm for head lettuce, salsify tops, endive, Chinese cabbage, 12 ppm for other leafy greens.
Leafy Vegetables*, Kale* Mustard Greens, Turnip Greens, Dandelion, Endive, Swiss Chard, Beet Greens, Spinach, Parsley, Lettuce, Collards, Chinese Cabbage	Flea beetles, leafhoppers, harlequin bug, Six spotted leafhopper (aster yellows vector), Corn earworm, armyworms, cabbage caterpillar, tarnished plant bugs and stink bugs		Day of harvest, Tolerance 5 ppm in ears. Tolerance 100 ppm on forage and fodder.
Corn*	Corn earworm, corn rootworm (larva), fall armyworm, sap beetles, Japanese beetle, flea beetles, leafhoppers		

For larvae in whorl (budworm) damage, apply to the plant. Repeat as necessary. For insects attacking stalks and ears, apply at 2 or 3-day intervals starting when first eggs appear, and continue until the stalks are dry. Three or more applications may be required depending on severity of infestation. Timing and frequency of applications may vary for other uses.

*When label directions for use are followed, forage, hay, and crop residues may be fed to meat or dairy animals.

ORNAMENTAL INSECT CONTROL - Good Way 5:50 (Dust) may be used for a wide range of ornamental plants, flowers, shrubs and small trees, such as rose, carnation, gladiolus, geranium, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. Good Way 5:50 (Dust) may be used for: Apply to the upper and lower surface of all foliage for complete coverage to control such pests as:

apple aphid, mealy bug, Japanese beetle, boxelder bug, leaf miner, scale, spider mite, willow leaf beetle, stink bug
bagworm, mimosa webworm, Japanese beetle, boxelder bug, leaf miner, scale, spider mite, willow leaf beetle, stink bug
birch leaf miner, oak leaf miner, Japanese beetle, boxelder bug, leaf miner, scale, spider mite, willow leaf beetle, stink bug
blister beetle, orange tortrix, leaf miner, Japanese beetle, boxelder bug, leaf miner, scale, spider mite, willow leaf beetle, stink bug
gypsy moth

NOTE Good Way 5:50 (Dust) is not recommended for use on plants which are recommended for use as a food source.

For protection of honeybees, do not apply to flowering plants which are visited by bees during the bloom period. When necessary, avoid spraying such periods, warn beekeepers when spraying, and do not spray directly on bees.

This product is highly toxic to bees exposed to direct treatment. Protective information may be obtained from your Cooperative Extension Service. Application of this product to the tree or shrub during the pollen shed period will seriously reduce bee population.

Injury to tender foliage may occur if applied to young plants. Avoid application to plants during the following treatment: AVOID CONTACT WITH EYES, EARS, WATER, UDDERS, BREASTS, MILK, FEEDING, OR DRINKING WATER. RECEPTACLE