

DANGER

Vapona Insecticide concentrate is poisonous if swallowed, inhaled or absorbed through skin or eyes. Do not get on skin or eyes or clothing. Do not breathe fume. Avoid prolonged exposure to fumes. Wear natural rubber gloves, protective clothing and goggles when handling the concentrate to prepare emulsions. Wear a mask or respirator of a type approved by the U.S. Bureau of Mines for DDVP, Vapona Insecticide. After use, keep a second person out of operational area.

In case of contact, wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash and decontaminate clothing with soap and water before reuse. DO NOT USE NEAR OPEN FLAME, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

ANTIDOTE: Atropine is the emergency antidote for vapona insecticide poisoning. Consult your physician about obtaining an adequate supply of 1/100 grain Atropine tablets for emergency use. Call a physician immediately in case of suspected poisoning. See "First Aid Treatment" Note to physician and first aid warnings and directions on side panels.

FIRST AID TREATMENT

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). Never give anything by mouth to an unconscious person.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the spill. Remove all contaminated clothing and wash skin with soap and running water. If the material gets into the eye, wash immediately with clean water at least 10 minutes.

Keep patient prone and quiet until a physician is available. Immediately if patient is not breathing.

If warning symptoms appear: Atropine, two 1/100 grain Atropine tablets, should be given immediately. Atropine should be given until warning symptoms disappear. See warning label for details.

NOTE TO PHYSICIAN

DANGER SYMPTOMS: Irritation of the respiratory tract, tightness in chest, blurred vision, dry mouth, dry nose, dry throat, sweating, nausea, vomiting, diarrhea and abdominal cramp.

TREATMENT: Atropine is the specific therapeutic antidote for the parasympathetic stimulation. If there are signs of parasympathetic stimulation, Atropine should be given at 10 minute intervals until the patient is free of symptoms. Complete response should be observed within 30 minutes.

Morphine is contraindicated. Clear intent to sedate or tranquilize patient is contraindicated. For 24 hours after exposure to this insecticide, the patient may show a marked increase in susceptibility to very small doses of any of the usual sedatives. Allow no further exposure until time for the patient to be operated on can be ascertained by the physician.

Distributed by
GOOD-WAY INSECTICIDE, INC.
Wheeling, Illinois

E.P.A. Reg. No. 2006-65

GOOD-WAY

VAPONA INSECTICIDE

CONCENTRATE

ANIMAL AND AREA SPRAY


KILLS: Face, Horn, Stable, House Flies, Mosquitoes and Certain

ACTIVE INGREDIENTS:	BY WEIGHT
2,2-Dichlorovinyl Dimethyl Phosphate (DDVP)	17.25%
Related Compounds	1.32%
Xylene	73.65%
INERT INGREDIENTS:	7.85%
TOTAL	100.00%

*Equivalent to 18.5% Vapona Insecticide Concentrate.
R. S. H. Chemical Co. Registered Trademark



CATTLE



THIS IS A CONCENTRATE—MUST BE DILUTED BEFORE USING

DIRECTIONS FOR USE: To make 1% water base spray, add 1 part Vapona Insecticide Concentrate to 16 parts water. To make 1.5% water base spray, add 1 part Vapona Insecticide Concentrate to 14 parts water. To make 1% Oil base fogging solution, add 1 part Vapona Insecticide to 20 parts fuel oil. Mix only enough dilute spray for one day's use.

RESTAURANTS, FOOD PROCESSING PLANTS AND CREAMERIES: Apply by train. Apply to areas around baseboards, cracks, walls, and window frames and floors. Do not apply to areas which will be contacted frequently by children, and do not allow children in treated areas until 24 hours after use. Do not contaminate food, water, food containers, or equipment. Do not remove fish bowls and aquaria before treatment.

Do not use in commercial food preparation areas or in other areas of food processing plant.

Not for use on storage bins or around the home.

FOGGING SOLUTION FOR ANIMAL BUILDINGS FOR CONTROL OF FLIES: Apply to barns, stables, stock pens, etc. Apply to walls, ceiling, and floor. Before applying, remove manure and other waste from the building. Wash walls and floor with water before application. Do not use in areas where animals are kept.

TO PROTECT DAIRY CATTLE AND BEEF ANIMALS: Apply to face, horns, and legs of animals. Use 1.5% water base spray. Apply spray daily at rate of 1 qt. per 100 lbs. of animal. Cover the animal with spray with particular attention to back, tail, and legs. Do not soak the skin or wet the hair. Mix only enough dilute spray for one application.

An 0.5% water base spray may be sprinkled on cow, horse, and pig for several hours. Apply with sprinking can at rate of 1 qt. per 100 lbs. of animal. Do not use in areas where animals are kept.

TO CONTROL FACE FLIES: Apply to face of animals. Use 1.5% water base spray. Apply to face of animals daily at rate of 1 qt. per 100 lbs. of animal. Do not use in areas where animals are kept.

DANGER IN USING: Do not use in areas where animals are kept. Do not use in areas where children are present. Do not use in areas where food is prepared. Do not use in areas where food is stored. Do not use in areas where food is consumed. Do not use in areas where food is distributed. Do not use in areas where food is sold. Do not use in areas where food is transported. Do not use in areas where food is stored. Do not use in areas where food is consumed. Do not use in areas where food is distributed. Do not use in areas where food is sold. Do not use in areas where food is transported.

NOT FOR USE IN OR AROUND THE HOME