


50 LBS. NET WEIGHT
VIGORTONE
INSECTICIDAL SALT NO. 151



For Continuous Self-Feeding of Cattle During the Summer for Control of Homflies and Cattle Grubs; for (Mixing in Feed) to Provide a 7-Day Feeding Program for Grub Control and as an Aid in the Reduction of Cattle Lice, or a 14-Day Feeding Program for Grub Control; for beef cattle and non-lactating dairy animals only.

Follow feeding directions carefully.

Pesticidally Active Ingredient:

Ronnel [0,0-dimethyl 0-(2, 4, 5-trichlorophenyl) phosphorothioate]	6.0%
Pesticidally Inert Ingredients	94.0%

INGREDIENTS

Salt, Diatomaceous Earth, Cane Molasses, Mineral Oil, Glyceryl Monostearate, Iron Oxide and Pyrogenic Silica.

WARNINGS

Use as a sole source of Ronnel.

Dairy cows should be treated early in their dry period but only if the dry period occurs between the end of the adult heel fly season and before grubs reach the back.

Do not feed to dairy animals that will produce milk for human consumption for 10 days prior to freshening. If dairy cows or heifers freshen during feeding program, or if insecticidal salt has not been withdrawn the required 10 days prior to freshening, milk must not be used for food for 10 days after the last feeding of this product.

Withdraw 10 days prior to slaughter.

Do not feed to sick animals or animals under stress, such as those just shipped, dehorned, castrated, or weaned within the last 3 weeks.

Ronnel is a cholinesterase inhibitor. Do not feed this product to animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides or chemicals. Consult a veterinarian at the first sign of adverse reaction. Atropine is an antidote.

CAUTION - KEEP OUT OF REACH OF CHILDREN

Wash hands with soap and water after handling. This material contains a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Use this product only as specified on this label. Do not reuse empty container. Destroy it by burning or burying in non-croplands away from water supplies.

EPA Est. 2011-IA-1; NB-1; OH-1 EPA Reg. No. 2011-3
 STORE IN A COOL, DRY PLACE

Rev. 5/74

Licensed under U.S. Patent No. 2,811,480

DIRECTIONS FOR CONTINUOUS SELF-FEEDING PROGRAM
VIGORTONE INSECTICIDAL SALT NO. 151

When to Feed: For hornfly control on pastured cattle start feeding Vigortone Insecticidal Salt No. 151 in the spring before flies appear. Continue feeding all summer and into the fall until cold weather restricts fly development. For grub control, feed continuously for a minimum of 75 days.

How Much to Feed: Recommended consumption of Vigortone Insecticidal Salt No. 151 containing 6% Ronnel is 0.2 lb. per 100 lbs. of body weight per head per month.

How to Feed: Feed Vigortone Insecticidal Salt No. 151 continuously free-choice to cattle in covered salt-feeder troughs or boxes. Provide one salt-feeder box or feeding space for each 15 to 20 head of cattle. Locate salt-feeder boxes and troughs near watering areas. To encourage uniform and normal consumption, put out only a 7-day supply at one time, and keep feeder boxes and troughs clean. For adequate insect control, each animal must eat its share. Small, late calves may require an additional 7 or 14-day feeding program after weaning for grub control.

WARNING

During continuous feeding program, do not allow cattle to run out of Vigortone Insecticidal Salt No. 151 and do not feed additional salt

DIRECTIONS FOR 7-DAY OR 14-DAY FEEDING PROGRAMS

Mix Vigortone Insecticidal Salt No. 151 with ground grain for top dressing, or mix with the complete ration to make an insecticidal feed. Feed continuously for 7 or 14 days and make certain that all cattle are eating. Adequate feeding space should be available for each animal. Make certain that insecticidal feed is evenly distributed on, or mixed into, the total ration. Provide a total of 0.2 lb. of Vigortone Insecticidal Salt No. 151 for each 100 lbs. body weight (to be fed over the 7 or 14-day feeding period).

NOTE: Insecticidal feed made from Vigortone Insecticidal Salt No. 151 should be used within 4 weeks of mixing.

When to Feed: After end of adult heel fly season and before grubs reach the back. Ronnel is not effective for grub control during later stages of life cycle, when grubs are in the backs of cattle. Reduction of cattle lice will be obtained when fed as directed for the 7-day program.

7-DAY TREATMENT

When to Feed:

- Southern area cattle - July through September
- Northern area cattle - August through October

How to Feed: Mix thoroughly 2 lbs. of Vigortone Insecticidal Salt No. 151 containing 6% Ronnel with 19 lbs. feed to produce an insecticidal feed. Prepare only enough for the 7-day feeding program. Feed 0.3 lb. of insecticidal feed per 100 lbs. body weight daily for 7 continuous days.

14-DAY TREATMENT

When to Feed:

- Southern area cattle - July through September
- Northern area cattle - August through **DECEMBER**

How to Feed: Mix thoroughly 1 lb. of Vigortone Insecticidal Salt No. 151 containing 6% Ronnel with 20 lbs. of feed to produce an insecticidal feed. Prepare only enough for the 14-day feeding program. Feed 0.3 lb. of insecticidal feed per 100 lbs. body weight daily for 14 continuous days.

VIGORTONE PRODUCTS COMPANY

A Division of Beatrice Foods Co.

Cedar Rapids, Ia. 52406 - Fremont, Neb. 68025 - Marion, Ohio 43302