

60 LBS 2.64 KG NET WEIGHT
VIGORTONE
BOVOTONE FC "007"
With Rabon[®] Oral Larvicide

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle

ACTIVE INGREDIENTS	BY WEIGHT
2-chloro-1,1,2,4,5-tetrahydrophthalazine-3,4-dithiophosphate	1.00%
INERT INGREDIENTS**	99.00%
TOTAL	100.00%

*Each pound contains 4.5g RABON[®]
 **Refers only to ingredients which are not larvicidal

EPA REG NO 20116

FEED INGREDIENTS

Dicalcium Phosphate, Monocalcium Phosphate, Calcium Carbonate, Salt, Yeast Culture, Corn Distillers Dried Grains with Solubles, Cane Molasses, Vitamin A Acetate, D Activated Animal Sterol, source of Vitamin D₃, Vitamin E Supplement, Magnesium Sulfate, Potassium Sulfate, Cobalt Carbonate, Calcium Iodate, Iron Carbonate, Manganese Oxide, Zinc Oxide, Copper Oxide, Sodium Benzoate, Iron Oxide, Selenium and Mineral Oil

GUARANTEED ANALYSIS

Calcium (Ca) maximum	15.6%
Calcium (Ca) minimum	13.0%
Phosphorus (P) minimum	7.00%
Salt (NaCl) maximum	22.00%
Salt (NaCl) minimum	19.00%
Iodine (I) minimum	0.007%
Vitamin A minimum	USP Units/lb 300,000
Vitamin D ₃ minimum	USP Units/lb 30,000
Vitamin E minimum	IU/lb 42

KEEP OUT OF REACH OF CHILDREN

CAUTION

**PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS**

CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal-Pesticide or pesticide that cannot be used according to label instructions must be disposed of according to federal, state and local procedures under the Resource Conservation and Recovery Act.

Container Disposal-Dispose of bags according to approved federal, state or local procedures under the Resource Conservation and Recovery Act.

Made in U.S.A.
VIGORTONE[®] AG PRODUCTS
 A Division of Beatrice Foods Co.
 Cedar Rapids, IA 52405 Fremont, NE 68025
 Marietta, OH 43302 Thomson, GA 30824

Printed 12/81

DIRECTIONS FOR USE - GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Allow free choice to cattle. Cattle should consume an average of 2.5 lb of the mixture per 100 lbs of body weight per month. This will supply the recommended average daily intake of 70 mg RABON[®] Oral Larvicide per hundred pounds of body weight. Animals should not be starved for minerals before initiation of the Vigortone Bovotone FC "007" with Rabon[®] Oral Larvicide program but OMIT all salt from other feed after the program is started. Do not feed other free choice sources of salt or mineral containing salt in any form. Allow one mineral feeder for each 10-15 head and place near watering and loafing areas. Put out no more than one month's supply at a time and protect from rain.

If consumption of Vigortone Bovotone FC "007" with Rabon[®] Oral Larvicide is above 2.5 lb per 100 lbs of body weight per month, reduce the number of feeding locations or reevaluate. If consumption is below 2.5 lb per 100 lbs of body weight per month, increase the number of feeding locations. Start feeding Vigortone Bovotone FC "007" with Rabon[®] Oral Larvicide early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall.

Vigortone Bovotone FC "007" with Rabon[®] Oral Larvicide may be fed up to slaughter and to lactating dairy cows.

Vigortone Bovotone FC "007" with Rabon[®] Oral Larvicide prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, Vigortone Bovotone FC "007" with Rabon[®] Oral Larvicide should be used in conjunction with other good management and sanitation practices.

MIXING DIRECTIONS FOR FINISHED FEEDS

The following table shows the correct amount of Vigortone Bovotone FC "007" with Rabon[®] Oral Larvicide to be mixed into a complete daily ration in order to provide the recommended average daily intake of 70 mg of Rabon[®] Oral Larvicide per hundred pounds of body weight.

Determine your cattle's average body weight, find this figure in the left column. Then determine the average consumption of your complete daily ration and find this figure in the top column. Cross reference and you will find amount of Vigortone Bovotone FC "007" with Rabon[®] Oral Larvicide to be mixed into your ration. Mixing time should be adequate to assure uniform dispersion. Optimum performance can be assured only if Vigortone Bovotone FC "007" with Rabon[®] Oral Larvicide is dispersed uniformly with the feedstuff at the recommended level. Common feed mixing equipment (i.e. vehicle mixers, horizontal blenders, mixer feeder trucks) may be used to prepare formulated feeds.

CONSUMPTION PER HEAD PER DAY OF FINISHED FEED (Pounds)

Wt. (LBS)	41	50	60	70	80	90	100	110	120	130	140	150	160	170	180	190	200	250	300
Amount	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	3000
Amount	1.0	1.25	1.5	1.75	2.0	2.25	2.5	2.75	3.0	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0	6.25	7.5
Weight	400	500	600	700	800	900	1000	1100	1200	1300	1400	1500	1600	1700	1800	1900	2000	2500	300