


5. **Disinfecting rubber goods:** Clean thoroughly, then soak in solution diluted 1 ounce to 1 gallon of water for 10 minutes.
6. **Cold disinfection of surgical instruments:** Clean thoroughly, rinse, and immerse in a solution of 1 ounce per gallon of water for 10 minutes. One tablespoonful of sodium nitrite should be added to each gallon of solution as a corrosion inhibitor.
7. **Sanitization of milk cans and miscellaneous dairy equipment:** Clean and rinse thoroughly. Then immerse in sanitizing solution diluted 1 ounce to 4 gallons of water. Milk cans and other equipment should be air dried. The same dilution is also recommended for flank and udder washing prior to milking.
8. **Sanitization of equipment in food processing plants, including meat plants:** Clean and rinse equipment thoroughly. Then, scrub with a solution of 1 ounce to 4 gallons of water (200 ppm quaternary). No rinsing is required. On surfaces difficult to clean completely (such as meat cutting boards) increase strength of sanitizing solution to 1 ounce to 1 gallon of water. (800 ppm quaternary). This must be followed by a rinse with potable water.

"Do not reuse empty container. Rinse thoroughly with water before discarding or returning drum to drum reconditioner"



NAMICO DISINFECTANT AND SANITIZER

Active Ingredients

n-alkyl (50% C ₁₄ , 40% C ₁₂ , 10% C ₁₆) dimethyl	
benzyl ammonium chloride	10%
Ethanol	2%
Inert Ingredients	88%

Phenol Coefficients

Salmonella typhosa	50
Staphylococcus aureus	75

E.P.A. Registration Number - 2021-18

DANGER

KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes severe eye and skin damage. Do not get in eyes, or skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID

External — In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before use.

Internal — If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN — Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock as well as oxygen and measures to support breathing manually or mechanically may be needed. If persistent, convulsions may be controlled by the cautious intravenous injection of a short-acting barbiturate drug.

ANOTHER NAMICO PRODUCT
NATIONAL MILLING & CHEMICAL COMPANY
Philadelphia 19127, Pa.
Contents — 1 Gallon

1. General disinfection of rooms, operating doors, etc.: Add 1 ounce of disinfectant to 1 gallon of water.
2. Barber shop and dental offices: soak combs, brushes, etc. in 1 ounce of disinfectant per gallon of water.
3. Sanitization of garbage pails, cutlery, etc.: Clean thoroughly, then immerse in 4 gallons of water (200 ppm quaternary). (Recommended by Federal Food, Drug, and Cosmetic Act)
4. Sanitizing clothing: Immerse in 4 gallons of water (200 ppm quaternary). Agitate for 10 minutes. Rinse thoroughly. Disinfectant in 2 ounces of disinfectant per gallon of water.

NAMICO DISINFECTANT AND SANITIZER

Active Ingredients

n-alkyl (50% C ₁₄ , 40% C ₁₂ , 10% C ₁₆) dimethyl	
benzyl ammonium chloride	10%
Ethanol	2%
Inert Ingredients	88%

Phenol Coefficients

Salmonella typhosa	50
Staphylococcus aureus	75

E.P.A. Registration Number - 2021-18

DANGER

KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes severe eye and skin damage. Do not get in eyes, or skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID

External — In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before use.

Internal — If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN — Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock as well as oxygen and measures to support breathing manually or mechanically may be needed. If persistent, convulsions may be controlled by the cautious intravenous injection of a short-acting barbiturate drug.

ANOTHER NAMICO PRODUCT

NATIONAL MILLING & CHEMICAL COMPANY
Philadelphia 19127, Pa.

Contents — 1 Gallon

USES

1. General **disinfection** of inanimate objects — hospital wards, sick-rooms, operation rooms, refrigerators, lavatories, urinals, hoppers, doors, etc.: Add 2½ ounces to 2 gallons of water, apply freely over surfaces to be disinfected.
2. Barber shop and beauty parlor equipment: Clean thoroughly, then soak combs, brushes, razors, etc., for 10 minutes in a solution of 1 ounce of **disinfectant** added to 1 gallon of water.
3. **Sanitization** of dishes, glassware, cooking utensils, silverware, bottles, garbage pails, cuspidors, etc. Clean with a suitable detergent, rinse thoroughly, then dip in a sanitizing rinse made with 1 ounce of product in 4 gallons of water, and air dry. **DO NOT USE A TOWEL.** (Recommended dilution should conform with local Health Department Regulations.)
4. **Sanitizing** clothes, towels, and linen — **COMMERCIAL LAUNDERING.** Agitate for 10 minutes in a solution containing 1 ounce of disinfectant in 2 gallons of water. **HOME LAUNDERING:** Use 1½ ounces of disinfectant to 1 gallon of water. Completely immerse fabric.

ED
6
cide.
le Act.
cide
1-18