EPA Form 8570-6 (Rev. 5-76)

SIGNATURE OF APPROVING OFFICIAL

DIRECTIONS FOR USE

It is a violation of Vederal Law to use this product in a manner inconsistent with its libeling.

APPLICATION INFORMATION

For use on hard, non-porous, manimate surfaces (glass, stainless steel, galvanized metal, porcelain, finofeum, tile, enameled and finished wood surfaces). Use on walls, floors, appliances, furnishings, and utensils.

Prepare fresh solutions daily. Do not reuse solutions. Discard solutions that are visibly soiled. Apply use solutions by flooding, immersion, swabbing, mopping or brushing to thoroughly wet all surfaces.

FOR DISINFECTION: In Hospitals, Nursing Homes, Food Plants, Commercial Freezers, Restrooms, Restaurants, end Veterinary Clinics, Farm Premises ara Poultry Houses. Use I oz. of this Disinfectant in 2 gallons of water for one-step cleaning and disinfecting light and moderately soiled surfaces. Heavily soiled surfaces should be cleaned and rinsed before disinfecting. For effective action, surfaces must remain wet for at least 10 minutes.

Food contact surfaces and all surfaces in edible food processing areas of Federally inspected meat and poultry plants must be rinsed with potable water before reuse when using the product at this concentration. Always carefully protect or remove food and packaging materials from the area before disinfecting.

For Farms and Poultry use, remove all animals, poultry and feed from premises. Remove all litter and manure from surfaces occupied or traversed by livestock, Empty all troughs, racks, and other feeding and watering equipment. Then, clean and disinfect as above. Ventilate buildings, coops, and other closed spaces. Do not house animals or poultry until treatment has been absorbed, set or dried.

disinfecting toilet boals and urinals: Remove gross soils from surfaces with bowl brush. Add I oz. of this Disinfectant to the bowl water. Use brush or swab to thoroughly clean bowl surfaces, especially under the rim. Flush commode or urinal several times and rinse bowl brush well.

SANITIZING OF NON-FOOD CONTACT SURFACES such as walls, floors, deck and counter tops, telephone receivers, door knobs, chairs, toilet seats, bathroom sinks, tubs, and shower stalls, in offices, hatels, incustrial and commercial buildings, and non-critical areas of hospitals and nursing homes: clean and sunitize in one operation by spraying or wiping a solution of 1 oz. of this Disinfectant in 3 gallons of water onto the sarfaces to be sanitized. Allow solution contact for 3 minutes before wiping clean or rinsing. If surfaces are f leavily soiled, preclean before sanitizing.

BEST AVAILABLE COPY

CHEMGO CHEMIDINE

ACCEPTED THE BEN GOT WIG COMMENTEQUIPMENT ! in EPA Letter Dallyshing, somproz. of this Di another suitabl

APR 0 & 1987 Disintectan

L-2_ութվուն cs. Hinder ti . ! Pangicide, and R is a control

Bactericide - Virucide meistered under EPA RORCHESSING
Food Plants-Livestock Buildings-Hospitals-Variation Clipics handlers, rins

NO RINSE FOOD PLANT SANITIZER POULTRY DRINKING WATER SANITIZER

*Kills Herpes Simplex Type II when used as directed for distinfection.

ACTIVE INGREDIENTS:

alpha-(P-nonyl phenyl)-omega-hydroxypoly (oxyethylene)-iodine complex:

(Providing 1.75% titratable iodine)

INERT INGREDIENTS: TOTAL INGREDIENTS:

KEEP OUT OF REACH OF CHILDREN DANGER

Practical Treatment: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or, if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory, shock, respiratory depression, and convultion may be needed.

FOR MEDICAL, INDUSTRIAL, COMMERCIAL, AND AGRICULTURAL USE ONLY.

A Product of

NET CONTENTS:

Cherago Corporation 10 Pennsylvania Avenue Malvern, PA 19355

I Gallen

EPA Reg. No. 2044-20 EPA Est. No. 2044-PA-L l-ozriper 5 g titratable lod

RETOR' **OPERATION** Disinfectant to control or and as neede reading of 17

SANITIZATI I ounce of water. Re control slim

SHOE BATI disease orga and hog ba disinfactant Scrape off: for 30 secon

> FOR CLE farrowing with a luke lo nolleg grain dry a

DEODORA it sanitize cations (ga solution of

STORAC container.

DISPOS. container

> Harmful damage (or on clo gloves w water af Cleffing