

DIRECT. : Read entire label before using

Hold container 6 to 10 inches from the animal and spray lightly over the entire body. Do not spray in eyes or on scrotum. For best penetration of spray to the skin, direct spray against the natural lay of the hair to cause fluffing of the coat.

Treat a 5 pound cat for 3-5 seconds or a 10 pound cat for 5-10 seconds. Apply enough to dampen hair and skin. Keep can in motion while spraying. Wet ticks thoroughly. Controls ticks for up to 1 week and fleas for up to 4 weeks. Treat a 15 pound dog for 10-20 seconds or a 24 pound dog for 20-30 seconds.

Repeat as necessary but not more often than once weekly.

MYCODEX FLEA & TICK SPRAY provides rapid kill of fleas on dogs and cats and of ticks on dogs. For more prolonged insect control, the animal's bedding should also be treated to help prevent re-infestation.

WARNING: Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing of spray mist and provide adequate ventilation of treated area. Contact with skin, eyes, or clothing should also be avoided. Wash thoroughly with soap and warm water after handling. Avoid contamination of food, utensils, and food preparation areas. If poisoning should occur, obtain prompt medical aid.

TO PHYSICIAN Large doses of active ingredient will result in cholinesterase depression. Atropine is antidotal. Do not give morphine. Watch for delayed pulmonary edema in cases of serious poisoning. In this case, use oxygen therapy and treat symptomatically.

NOTICE When used in accordance with these directions MYCODEX Flea & Tick Spray is effective. However, because of the wide range of conditions under which the product may be used even though these directions are followed, the user accepts all risks and responsibilities.

WARNING: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures over 130° F may cause bursting. Never throw container into fire or incinerator.

EPA Reg. No. 2097-9
Est. No. 10807-GA-1

W75

CONTENTS:
Active Ingredient
Inert Ingredients

0 Isopropoxyphenoxy Methyl Carbamate 0.25
99.75
U.S. Patent No. 3,111,539

For Veterinary Use Only

KEEP IN COOL, DRY PLACE

CAUTION: Keep Out of Reach of Children.

See other precautions on the back/side panel.

NET WT. 12 OZ.

Beecham
Laboratories