

**NACO**

**NACO**



**DIRECTIONS FOR USE**

Sulphur Smoke Dusting Sulphur is formulated for use in dust mixtures or alone as a dust when a 325 mesh dust is required. For best results apply when wind is calm. Dust often enough to keep foliage and fruit covered with a light uniform coating. Consult with your State Agricultural Experiment Station or your State Agricultural Extension Service for additional information consistent with this label cautions and directions.

**APPLES:** To control Scab and powdery mildew apply 30 to 40 lbs. per acre in the green tip to pink bud stage. Use on sulphur tolerant varieties only.

**PEACH AND NECTARINE:** To control Brown rot, coryneum blight, rust, peach silver mite and powdery mildew apply 50 to 60 lbs. per acre from popcorn stage to harvest. Repeat at weekly intervals.

**PLUM AND PRUNE:** To control Brown rot, leaf spot, and prune rust apply 35 to 50 lbs. per acre starting 3 weeks before harvest and continuing at weekly intervals.

**BEANS AND PEAS:** To control powdery mildew, rust, and aphid apply 25 to 30 lbs. per acre. Start application as soon as the first true leaves form and continue at weekly intervals until late podding. May cause injury to some varieties of beans.

# SULPHUR SMOKE<sup>®</sup>

## DUSTING SULPHUR

**ACTIVE INGREDIENTS:**

Sulphur*	93.00%
INERT INGREDIENTS	7.00%
<b>TOTAL</b>	<b>100.00%</b>

\*Mesh Test not less than 90% through 325 mesh.  
† Sulphur Smoke Reg. trade name U. S. Patent Office

### CAUTION!

#### KEEP OUT OF REACH OF CHILDREN

Do not use sulphur within 3 to 6 weeks of an oil spray. Do not use sulphur on apples, apricots, Anjou and Comice pears, cranberries, cucumbers, cucumbers, squash and some varieties of melons, filberts, spinach, tung trees, and walnuts when temperatures exceed or are likely to exceed 90° F.

W. R. GRACE & Co  
AGRICULTURAL CHEMICAL  
MEMPHIS, TENNESSEE

NET WEIGHT 50 LBS.

USDA REG. NO. 2124-62

**DIRECTIONS Continued**

**COTTON:** To control Atlantic mite, and cotton rust apply 25 to 30 lbs. per acre. Start application as soon as the first true leaves form and continue at weekly intervals until blossom time.

**MELON (Honeydew and sulphur-resistant melons only):** To control Powdery mildew and spider mites apply 20 to 25 lbs. per acre. Repeat at weekly intervals.

**PEANUTS:** To control Rust and leaf spot apply 15 to 30 lbs. per acre. Repeat at 7 to 10 day intervals.

**CITRUS:** To control Sc-splotted and citrus rust mites apply 80 lbs. per acre (1 1/2 to 1 1/2 lbs. per tree, depending on size, for citrus rust mites in Florida.) To control Citrus flat and Yuma spider mites apply 100 lbs. per acre. Do not apply within 3 to 6 weeks of an oil spray.

**NOTICE:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material in accordance with directions given hereon.

*Handwritten:* Rec 5/19/71  
2124-62