

UNDER THE FEDERAL INSECTICIDE ACT

hands, arms and face thoroughly with soap and warm water before eating or smoking.

Wash all contaminated clothing with soap and hot water before re-use.

SYMPTOMS OF POISONING: A sense of "tightness" in the chest. Sweating.

Contracted pupils. Stomach pains. Vomiting and diarrhea.

TREATMENT: In case of poisoning call a physician immediately. Have patient lie down and keep quiet. If swallowed, induce vomiting by giving a tablespoonful of salt in a glass of warm water. Administer milk or water freely and again induce vomiting until fluid is clear. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash with flowing water for at least 15 minutes.

ANTIDOTE: Administer atropine sulfate in large therapeutic doses. 2-PAM is also antidotal and may be administered in conjunction with atropine. Repeat as necessary to the point of tolerance.

TO PHYSICIAN: Compound inhibits cholinesterase resulting in stimulation of the central nervous system, the parasympathetic nervous system, and the somatic motor nerves. Do not give morphine. Watch for pulmonary edema, which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically. Do not contaminate feed and foodstuffs.

Not for use or storage in or around the home.

This product is poisonous to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on the label. Do not apply to water drainage areas where runoff or flooding will contaminate ponds, lakes or streams. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

ACCEPTED

The principle active ingredient of this product is patented under U.S. Patent No. 2,759,010 and Canadian Patent No. 518,967 of Chemagro Corporation, Kansas City, Missouri, and the purchase includes a limited license under said patents to use or reformulate the product only in the manner provided by the recommended applications on the present label. The purchaser obligates itself to notify Chemagro Corporation of any proposed use or reformulation of such licensed product not provided by said recommended applications, and to obtain Chemagro's approval in writing before making any such use or reformulation of the licensed product. Any use, or reformulation intended for use, of the licensed product in the home and garden field is specifically prohibited by said license.


RECOMMENDED DI-SYSTON APPLICATIONS

IMPORTANT: When used on Tomatoes, or Beans at the maximum recommended rates under adverse conditions such as extremely cool and wet or extremely dry weather, DI-SYSTON may cause some delay in emergence, stunting of seedlings, or reduction of stand. Such damage may be more pronounced in light sandy soils. Plant injury may also occur or may be enhanced when certain pre-emergent herbicides are used in conjunction with DI-SYSTON.

DIRECTIONS FOR USE

DI-SYSTON is a systemic insecticide entering the plant by absorption through the roots. The insecticide placed near the roots is translocated to all parts of the plant and kills sucking insects feeding on the plant. The active ingredient is released slowly giving long protection during the plant's susceptible early life.

3134-129



VEGETABLES:

BEANS: To control Aphids, Leafhoppers, Mexican bean beetle (Larvae), Mites and Thrips, apply at the rate of 250 - 500 lbs. (1 to 2 lbs. actual DI-SYSTON) per acre. For Snap or Green Lima Beans, apply specified dosage per acre in a band on each side of the seed furrow at planting time only. For Dry Beans apply specified dosage per acre as above at planting time or as a side-dressing on each side of the row after plants become established but not within 60 days of harvest. Do not apply directly to seed. Do not apply more than once per season. Read "IMPORTANT" statement under "Recommended DI-SYSTON Applications" for additional information before use on this crop.

BRUSSELS SPROUTS & CAULIFLOWER: To control Aphids, Flea beetles, Leafhoppers, Mites and Thrips on Brussels Sprouts and Aphids, Flea beetles and Leafhoppers on Cauliflower apply 250 lbs. (1 lb. actual DI-SYSTON) per acre. Apply specified dosage per acre in a band on each side of the seed furrow at planting or as a side-dressing after plants become established. Do not apply directly to seed. Do not apply more than twice per season, nor within 30 days of harvest for Brussels Sprouts, or 40 days of harvest for Cauliflower. Allow a minimum of 21 days between applications.

BROCCOLI AND CABBAGE: To control Aphids, Flea beetles, Leafhoppers on Broccoli and Aphids, Flea beetles on Cabbage apply 250 lbs. (1 lb. actual DI-SYSTON) per acre. To control Root aphid on Cabbage apply 375 lbs. (1 1/2 lbs.

RECEIVED

actual DI-SYSTON) per acre. Apply specified dosage per acre in a band on each side of the seed furrow at planting or as a side-dressing after plants become established. Do not apply directly to seed. Do not apply more than once per season. Do not use within 14 days of harvest for Broccoli or 42 days of harvest for Cabbage.

POTATOES: To control Aphids, Flea beetles (except Tuber flea beetles), Leafhoppers, Potato psyllid, Colorado potato beetle (early season reduction), Southern potato wireworm, apply at the rate of 500 - 750 lbs. (2 to 3 lbs. actual DI-SYSTON) per acre. Apply specified dosage per acre in the furrow or in a band on each side of the seed furrow at planting or as a side-dressing fertilizer after plants become established. Do not make more than two (2) applications per season. Do not apply within 15 days of harvest. Higher rate to be used in organic soils.

SPINACH: To control Aphids, Leafhoppers, Mites and Thrips on Spinach, apply 250 lbs. (1 lb. actual DI-SYSTON) per acre. Apply specified dosage per acre in a band on each side of the seed furrow at planting time only. Do not apply directly on the seed.

TOMATOES: To control Aphids, Flea beetles, Leafhoppers, Leaf miners, Colorado potato beetle (early season reduction) and Mites on Tomatoes, apply 250 to 750 lbs. (1 to 3 lbs. actual DI-SYSTON) per acre. Apply maximum specified dosage per acre in a band on each side of the seed furrow at planting time only. Do not apply directly to seed. Where 2 treatments are required, use only 250 - 500 lbs. per acre per treatment applied in a band on each side of the seed furrow at planting and in

ACCEPTED
11/11/11

a side-dressing after plants become established. Allow a minimum of 21 days between applications. Do not apply within 30 days of harvest. Read "IMPORTANT" statement under "Recommended DI-SYSTON Applications" for additional information before use on this crop.

FIELD CROPS:

ALFALFA: To control Aphids and Mites apply at the rate of 250 lbs. (1 lb. actual DI-SYSTON) per acre. Apply specified dosage per acre by broadcast as needed. Allow a minimum time of 28 days between applications.

BARLEY AND OATS: To control Aphids and Mites, apply at the rate of 250 lbs. (1 lb. actual DI-SYSTON) per acre. Apply specified dosage per acre. Apply specified dosage per acre by drilling or broadcast at planting time, or as a broadcast after emergence. Do not apply more than twice per season nor within 60 days of harvest. Allow a minimum of 21 days between applications. Do not graze or cut for forage within 30 days of treatment.

CORN: To control Rootworms, apply at the rate of 250 lbs. (1 lb. actual DI-SYSTON) per acre. Apply specified dosage per acre at planting time in the starter fertilizer. Do not harvest treated corn for food, feed or ensilage within 100 days of treatment.

WHEAT: (Fall) To control Hessian Fly, apply at the rate of 250 lbs. (1 lb. actual DI-SYSTON) per acre. Apply specified dosage per acre by drilling or broadcast at planting time in the fall.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended

and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

W. R. GRACE & CO.
AGRICULTURAL CHEMICALS
MEMPHIS, TENNESSEE