



CORONA

ACTIVE INGREDIENTS:	
Lead Arsenite (PbHgAsO <sub>4</sub> ) .....	98.6%
INERT INGREDIENTS .....	2.0%
Arsenic expressed as metallic	21.14%
Arsenic in water soluble form expressed as metallic	
Not more than	0.35%
Lead expressed as metallic	58.6%

(16-point)

DANGER!

~~DO NOT DRINK OR EAT DURING USE.~~

## POISON

**SYMPTOMS:** Give a full teaspoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Then give two tablespoonsfuls of spirit. Gums on teeth of the victim in water or a mixture of milk and water. Have victim lie down and spit. **DO NOT INDUCE VOMITING.** **DO NOT INDUCE SPONTANEOUS DEFECATION.** **AVOID PRACTICING** **ARTIFICIAL RESPIRATION.** **KEEP FROM EYES AND MOUTH.** **AVOID PRACTICING** **ARTIFICIAL RESPIRATION.**

Ingestion: induce emesis and gastric lavage.

Do not apply to water. Keep out of any body of water. Apply directly to only areas specified on this label. To protect bees, avoid application of this product directly on blossoms. Wash spray equipment thoroughly after use. If 100% of wash water is sprayed, it will not be a hazard to humans and animals.

### DIRECTIONS

For control of cockroaches, ants, beetles, and other insects, apply to infested areas and surfaces. On fruit and vegetables, mix one quart of water with one-half cup of insecticide. Spray until surface is moistened. Do not spray with the nozzle pointed at人.

Caution: Effects may be delayed and may not be apparent immediately.

2(0P3)

-2-

2. Cherry: Control of Black Cherry Borer, Cherry Leafworm, and Cherry Fly. Make 2 to 3 applications at 7 to 10 day intervals, beginning at petal fall or shuck split. Make no application later than 5 days before harvest. Remove excess residue at harvest.

CHERRIES: For control of Plum Curculio use 2 to 3 lbs. per 100 gals. Make 2 to 3 applications at 8 to 10 day intervals, beginning at petal fall or shuck split. For Cherry Fruit Flies use 2 lbs. lead arsenate plus 3 lbs. hydrated lime per 100 gals. water; apply when first flies emerge; make second application 10 days later. Do not apply later than 14 days before harvest on cherries to be processed, nor 7 days on cherries for the market.

\*On the 1st and 2nd applications, water should be reduced to 1 lb. per 100 gals. water.

LEAFWORM: Control of Leafworm, Hornworm and Cabbage Looper. Use 2 to 3 lbs. lead arsenate, lime, hydrated lime or other infuser per 100 gals. water. Apply 1 to 2 lbs. of spray per acre, depending on size of plants. Make 2 to 3 applications; repeat as needed for control. Remove excess residue at harvest.

CHERRY FLY: For control of the Cherry Fly, use the following. Make 2 applications, one at petal fall, the second 7 to 10 days after the first. Consider use of "Derris" or "Garbanzo" insecticide if the above treatment does not give satisfactory results. For additional information concerning the use of Derris and Garbanzo see the "Derris and Garbanzo" section.

375-1000

100-1000

100-1000

3/0f3

W.R.G.: W.R. Grace & Co. Inc. Timing and method of application, weather and crop condition, mixing with other chemicals and especially temperature, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

W. R. GRACE & CO.  
AGRICULTURAL CHEMICALS

NET WEIGHT POUNDS