


NACO

NACO



USDA REG. NO. 2124-466

MICRONIZED WETTABLE SULFUR

DIRECTIONS. Continued

combine 1 gal. of lime sulfur solution (or 4 lbs. dry lime sulfur) with 3 to 4 lbs. of Micronized Wettable Sulfur in 100 gals. of water.

ON CALYX AND COVER APPLICATIONS use 4 to 6 lbs. of Micronized Wettable Sulfur in 100 gals. of water depending upon degree of scab infection and weather conditions. As an alternate method, NACO Micronized Wettable Sulfur may be used at 3 to 5 lbs. per 100 gals. with $\frac{3}{4}$ lb. of Coromate. This combination is very effective in controlling scab on McIntosh during periods of heavy infection.

For PEACHES (Brown Rot and Peach Scab): Use 3 to 4 lbs. of Micronized Wettable Sulfur in 100 gals. of water in spray applications for shuckfall; then two weeks and four weeks after and in any necessary additional applications. When Arsenate of Lead is to be used with sulfur on peaches or other stone fruits, use hydrated lime in an amount equal to the Arsenate of Lead. When Zinc Sulphate is used with this mixture as a peach spray, add the ingredients to the contents of the spray tank — with

ACTIVE INGREDIENTS		
Sulfur	95.00%	
INERT INGREDIENTS	5.00%	
TOTAL	100.00%	

agitator running in the following order: Micronized Wettable Sulfur, Zinc Sulphate, Hydrated Lime, Arsenate of Lead.

CAUTION: For the maximum control of Apple Scab and Peach Brown Rot, the sulfur must be uniformly distributed over the plant surfaces to be protected before spore discharge. Do not apply sulfur at temperatures greater than 85 degrees F. or within two weeks of an oil application.

Consult your State Agricultural Experiment Station or your State Agricultural Extension Service for additional information consistent with this label cautions and directions.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

CAUTION!
KEEP OUT OF REACH OF CHILDREN

DIRECTIONS FOR USE

When tank is approximately two-thirds filled with water, sift in the Micronized Sulfur with the agitator running.

FOR APPLES (Scab): On PREPINK AND PINK SPRAYS use 6 to 8 lbs. of Micronized Wettable Sulfur in 100 gals. of water of

W. R. GRACE & CO.
AGRICULTURAL CHEMICALS
MEMPHIS, TENNESSEE

Amarillo, Tex. • Baltimore, Md. • Cary, N.C. • Charleston, S.C. • Columbus, Ohio • Ft. Pierce, Fla. • Henry, Ill. • Joplin, Mo. • Lansing, Mich. • Nashville, Tenn. • New Albany, Ind. • Odessa, Tex. • Perry, Iowa • Portland, Oregon.

NET WEIGHT 30 LBS.