

GRACE GRACE GRACE GRACE GRACE GRACE

GRACE GRACE GRACE GRACE GRACE GRACE

BEFORE USING
ANY PESTICIDE

STOP

READ THE LABEL

NACO

USDA Reg. No. 2124-480

PHALTAN® FOLPET 50 WETTABLE

FUNGICIDE ACCEPTED

<p>ACTIVE INGREDIENT: Folpet*</p> <p>INERT INGREDIENTS</p> <p>TOTAL</p> <p>• Phaltan is a registered trademark for Folpet Fungicide * N-(1-chloromethylthio) phthalimide U.S. Patent No. 2,553,770 — 2,553,771</p>	<p>50%</p> <p>50%</p> <p>100%</p> <p style="font-size: 2em; opacity: 0.5;">31</p>
---	---

CAUTION!
KEEP OUT OF REACH OF CHILDREN
(See additional cautions below)

AVOID INHALATION OF DUST OR SPRAY MIST. AVOID PROLONGED OR REPEATED CONTACT WITH SKIN. AVOID CONTAMINATION OF FEED AND FOODSTUFFS.
Do not use on Sweet Cherries as injury may occur. Store In Cool, Dry Place, Protect from Excessive Heat. Burn Bag Immediately When Empty. Stay Out of Smoke.
This product is poisonous to fish. Keep out of any body of water.
Use strictly in accordance with cautions, warnings, and directions on this label. Consult your State Agricultural Experiment Station or your State Agricultural Extension Service for additional information consistent with label directions and cautions.
USE DESIGNATED AMOUNT IN 100 GALLONS WATER UNLESS OTHERWISE SPECIFIED.

DIRECTIONS FOR USE

APPLES, CRABAPPLES: Bitter Rot, Botryosphera, Alternaria Spot, Sooty Blotch, Black Rot, Fly Speck and Scab—1 1/2 to 2 lbs. make first application at delayed dormant and repeat at 7 to 14 day intervals until harvest. Caution: East—used during the critical period just before bloom until 30 days after petal fall may cause russetting to sensitive varieties. West—Do not use between bloom and July 1 to avoid injury.

AVOCADOS: Scab—3 lbs. Make applications at 2 to 3 week intervals as needed.

BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, RASPBERRIES: Fruit Rot, Gray Mold, Spur Blight—2 lbs. Apply 250 gals. per acre. Make first application when berries have set and repeat at 7 day intervals until harvest.

BLUEBERRIES, HUCKLEBERRIES: Gray Mold, Powdery Mildew, Leaf Spot, Gloeosporium—2 lbs. Apply 250 gals. per acre. Make first application before first infection. Repeat at 7 day intervals until harvest.

CANTALOUPE, MUSKMELONS, HONEYDEW MELONS, PUMPKINS, WATERMELONS, WINTER SQUASH: Downy Mildew, Powdery Mildew, Anthracnose—2 to 4 lbs. Apply up to 200 gals. per acre depending on size of vines. Make first application when first true leaves appear and repeat at weekly intervals until crops is harvested.

CELERY: Early Blight, Late Blight, Cercospora Leaf Spot—2 lbs. Apply after plants are set in field and established. Repeat at 7 day intervals. Do not apply within 7 days of harvest.

CHERRIES (SOUP ONLY): Leaf Spot and Brown Rot—2 lbs. Apply at pre-bloom and popcorn stage. Repeat in bloom and at 7 to 14 day intervals making last application to foliage post-harvest.

CITRUS (Oranges, Grapefruit, Lemon, Lime, Tangerine, Tangerines): Melanose Scab Greasy Spot—2 lbs. Apply dormant, 2 1/2 petal fall, 2 weeks later and on the fall flush in August and September.

CRANBERRIES: Storage Rot—9 lbs. per acre. Apply during bloom and repeat at 10—14 day intervals. Do not apply within 30 days of harvest.

CUCUMBERS, SUMMER SQUASH: Anthracnose, Downy Mildew, Powdery Mildew—7 to 4 lbs. Apply to 200 gals. per acre depending upon size of vines. Make first application when first true leaves appear and repeat at weekly intervals until crop is harvested.

GRAPES: Downy Mildew, Powdery Mildew, Black Rot, Dead Arm—2 lbs. Apply 250 to 300 gals. per acre. Dead Arm—East—Make first application when new growth is 4 to 6 inches long, repeat when growth is 8 to 10 inches long. West—first application bud break to 1 inch shoot length. Second application 2 weeks later or 4 to 8 inch shoot length. Black Rot, Downy Mildew—Make first application just before bloom, repeat immediately after bloom. Depending upon the susceptibility of the grape variety from 1 to 3 more sprays may be needed at 7 to 10 day intervals.
Powdery Mildew—East only—Make first application 2 to 3 weeks after bloom and repeat at 2 week intervals as needed.
Note: During extended periods of hot dry weather, foliage injury to several varieties of grapes may occur.

GOOSEBERRIES, CURRANTS: Leaf Spot—4 lbs. Apply 150 gals. per acre. Make first application before first infection. Repeat at 7 day intervals until crop is harvested.

LETTUCE: Downy Mildew—2 lbs. Apply up to 200 gals. per acre. Make first application when plants are established in field. Repeat at 7 day intervals until harvest.

ONIONS, GARLIC, LEKES, SHALLOTS: Purple Blotch and Downy Mildew—Use 4 lbs. Apply 200 gals. per acre. Make first application before first infection. Repeat at weekly intervals until tops begin to dry before harvest.

STRAWBERRIES: Leaf Spot, Fruit Rot—2 lbs. Apply up to 200 gals. per acre. Make first application before bloom and repeat at 7 day intervals until after harvest.

TOMATOES: Anthracnose—4 lbs. Use up to 200 gals. per acre. Make first application during first bloom and repeat at 7 to 10 day intervals until fruit is harvested.

ROSES: Black Spot, Powdery Mildew, Rust—2 lbs. per 100 gals. Make first application when young leaves first appear. Repeat at 7 to 10 day intervals or as necessary throughout the season. Make thorough application.

CHRYSANTHEMUMS, IRIS: Septoria Leafspot, Didymella Leafspot—2 lbs. per 100 gals. Start treatments immediately after planting. Repeat at 7 to 10 day intervals or as needed throughout the season.

CARNATIONS: Rust, Alternaria Leafspot—2 lbs. per 100 gals. Begin applications 2 weeks after planting and continue bi-weekly application for 3 to 4 applications.

ZINNIAS, MARIGOLDS, CHRYSANTHEMUMS, ASTERS, PHLOX, SNAPDRAGONS: Alternaria Leafspot, Powdery Mildew, Anthracnose—2 lbs. per 100 gals. Treat at least weekly during periods of rapid new growth and severe infection, and twice weekly during periods of rapid new growth and severe infection caused by continuous rains or high humidity.

POINSETTIA (GREENHOUSE): Pythium Root Rot—2 lbs. per 100 gals. water plus spreader and use 1 pint per 8 inch pan. Make first application at time of panning. Repeat in 10 days.

AZALEA CUTTINGS: Damp-off 1 1/2 to 2 lbs. per 100 gals. Allow to soak for 15 to 30 minutes before placing in rooting media.

COMPATIBILITY: NACO FOLPET 50 Wettable may be combined with most commonly used insecticides, adjuvants and fungicides with the exception of strongly alkaline material such as hydrated lime, which reduces the fungicidal activity of NACO FOLPET 50 Wettable. NACO FOLPET 50 Wettable should not be used in combination with or closely following oil sprays. Do not use with TEPH formulations when a copper fungicide mixture is being applied.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are ever changing. The seller, Buyer assumes all risk of use, storage and handling of this material. Not in strict accordance with directions given here-with.

W. R. GRACE & CO.
AGRICULTURAL CHEMICALS
MEMPHIS, TENNESSEE

Amarillo, Tex. • Baltimore, Md. • Cary, N.C. • Charleston, S.C. • Columbus, Ohio • Ft. Pierce, Fla.
Henry, Ill. • Joplin, Mo. • Lansing, Mich. • Nashville, Tenn. • New Albany, Ind. • Odessa, Tex.
Perry, Iowa • Portland, Oregon

GRACE GRACE GRACE GRACE GRACE GRACE

GRACE GRACE GRACE GRACE GRACE GRACE

NET WEIGHT LBS.

GRACE GRACE GRACE GRACE GRACE GRACE