

NACO

USDA Reg. No. 2124 585



**LAWN AND ORNAMENTAL
TURF TREATER
GRANULES**

ACTIVE INGREDIENTS:

Heptachlor (Heptachlorotetrahydro-4, 7-methanoindene) 5.00%
Related Compounds 1.76%

INERT INGREDIENTS

TOTAL 93.24%
100.00%

CAUTION!

KEEP OUT OF REACH OF CHILDREN

(See additional cautions and directions below)

HARMFUL IF SWALLOWED. AVOID INHALATION OF DUST.

Wash hands, arms and face with soap and water after using.

Avoid contamination of feed and foodstuffs.

ANTIDOTE: External — In case of spillage on skin, wash promptly with soap and water.

Internal — Emetic of 1 tablespoonful mustard in a tumbler of water. Call a Physician.

During commercial or prolonged exposure in mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type found suitable by the U. S. Department of Agriculture for protection against Heptachlor.

This product is toxic to fish and wildlife. Keep out of lakes, streams, and ponds. Birds feeding on treated areas may be killed. Apply this product only as specified on this label. Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning (Keep out of smoke).

DIRECTIONS FOR USE

Naco Lawn and Turf Treater Granules (containing 5% Heptachlor) is for use as a soil application only on lawns, park areas, golf courses, cemeteries and ornamental nurseries for control of certain insect pests. Apply with a combination drill or fertilizer spreader. Consult your State Agricultural Experiment Station or your State Agricultural Extension Service for additional information on usage of this product when consistent with this label cautions and directions.

LAWN AND ORNAMENTAL TURF: For control of fire ants apply 9 oz. per 1000 sq. ft. Wash thoroughly after application. The best time for treatment is from late fall to early spring. Do not apply to areas planted to food crops.

This product may be applied as a mound treatment (though not as effective as treating the entire area) where the infestation is light or for temporary control. Tear down the mound with a rake or hoe and apply 2 to 4 cupfuls per mound applied to the ground within a 3 foot area on all sides. Do not apply to areas planted to food crops.

For control of Chiggers, fleas and ticks apply 1/2 lb. per 1000 square feet (20 lbs. per acre).

Apply to grass, turf or ornamentals. Treat only infested areas. One treatment made when infestation is first noticed will usually last all season.

For control of Asiatic Garden Beetle Larvae, European Chafer Larvae, Japanese Beetle Larvae and White-Fringed Beetle Larvae apply 1 lb. per 1000 sq. ft. Apply to soil surface before planting and wash into the top few inches of the soil.

Do not apply to areas planted to food crops.

For control of Wireworms apply 1 lb. per 1000 sq. ft. Work into the soil.

CAUTION: Keep children and pets off treated areas until material is washed into the soil and area thoroughly dried.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

W. R. GRACE & CO.

AGRICULTURAL CHEMICALS

MEMPHIS, TENN.

Baltimore, Md. • Cary, N.C. • Charleston, S.C. • Columbus, Ohio • Ft. Pierce, Fla. • Henry, Ill.
Joplin, Mo. • Lansing, Mich. • Nashville, Tenn. • New Albany, Ind. • Amarillo, Tex.
Perry, Iowa • Portland, Oregon

NET WEIGHT 5 LBS.

Lot 100 - Rev. 2

2124-585