



SEVIN*-MANEB 5-1.5 EC

ACTIVE INGREDIENTS:	
Carbaryl (1-naphthyl N-methylcarbamate)	5.0%
Maneb (Manranous ethylene bisdithiocarbamate)	4.8%
INERT INGREDIENTS	90.2%
TOTAL	100.0%

*Trademark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate
 U. S. Patent Nos. 2,903,478; 3,009,555

(18-point)
 (12-point)

CAUTION!
KEEP OUT OF REACH OF CHILDREN
 (See side panel for additional cautions and directions)

HARMFUL IF INHALED OR SWALLOWED. DO NOT TAKE INTERNALLY. DO NOT BREATH DUST.
 SKIN CONTACT MAY BE HARMFUL. AVOID UNNECESSARY CONTACT. WASH HANDS BEFORE EATING.
 TAKE SHOWER OR BATH AFTER WORK. WEAR RUBBER GLOVES AND CLOTHING. CHANGE TO
 CLEAN CLOTHING DAILY.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor.
 Atropine is antidotal.

Avoid contamination of food stuffs. Avoid contamination of feed, feed receptacles,
 and water receptacles.

For protection of honeybees: Avoid use if bees are present near application area. Honeybees
 are visiting the crop or adjacent areas. When necessary to use in such
 periods, warn beekeepers well in advance to locate hives at a safe distance (be-
 yond the flight range) until one week after application.

DIRECTIONS FOR USE

Apply when the wind is calm at the rate indicated below. Be sure to cover all
 sides of the leaves with a light covering of dust. Repeat applications if needed
 occur within 7 days after application. Do not apply during flowering.

Do not graze treated fields or feed trash to dairy or meat animals.

TOMATOES: Apply 25 - 40 lbs. per acre.

Diseases - Early blight, late blight, Septoria leaf spot, gray leaf spot (stemphylium) and anthracnose - start applications when fruit of first clusters is well formed; repeat at 7 - 10 day intervals. (Do not use on tender young seedlings in greenhouses as injury may occur.) When weather conditions favor disease, more frequent applications may be necessary starting when plants are set in the field.

Insects - Tomato fruitworm, fall armyworm, flea beetles, tomato hornworm, European corn borer, and stink bugs. Apply at 7 - 10 day intervals when insects or their damage appear. The higher rate of application is necessary for hornworms, corn borer and stink bugs.

POTATOES: Apply 25 - 40 lbs. per acre.

Diseases - Early blight and late blight. Start applications as soon as the plants are 3 - 4 inches high and continue at 7 - 10 day intervals as long as blight threatens. Under severe disease conditions, use the shorter interval or higher dosage.

Insects - Colorado potato beetle, leafhopper, and flea beetle. Apply at 7 - 10 day intervals when insects or their damage appear.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

W. R. GRACE & CO.
AGRICULTURAL PRODUCTS DIVISION
BALTIMORE, MARYLAND

NET WEIGHT POUNDS

STO. 19, 1962
2124628