

ACCEPTED

2124-648

5-8-74

U.S. DEPARTMENT OF AGRICULTURE
NATIONAL INSECT MEDICATION BOARD
WASHINGTON, D.C. 20250**NACO**

EPA REG. NO. 2124-648

CHLORDANE 10 DUST

ACTIVE INGREDIENT:

Technical Chlordane*

INERT INGREDIENTS

TOTAL

10.0%

90.0%

100.0%

*Equivalent to 6.00% Octachloro-4, 7-methanotetrahydroindane and 4.0% Related Compounds.

CAUTION!

KEEP OUT OF REACH OF CHILDREN**HARMFUL IF SWALLOWED. CONTACT WITH SKIN CAN CAUSE TOXIC SYMPTOMS. AVOID BREATHING DUST.**
In case of contact with skin, wash with soap and water.**ANTIDOTE:** External--In case of spillage on skin, wash promptly with soap and water.

Internal--Emetic of 1 tablespoonful of mustard in tumbler of water. Call a physician.

Avoid contamination of feed and foodstuffs.

This product is toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Birds and other wildlife in treated areas may be harmed. Birds feeding on treated areas may be killed. Do not apply in any manner not specified on this label. Do not apply when weather conditions favor drift of dust from areas treated. Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by burying with wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

Consult with your State Agricultural Experiment Station or your State Agricultural Extension Service for additional information consistent with this label cautions and directions.

HOME LAWNS AND GOLF COURSES: To control Ants, Chiggers (red bugs), Chinch bugs, Fall armyworms, Green June beetle grubs, Mole crickets, Spittle bugs, and Wireworms use 100 lbs. of this product per acre. May be scattered evenly over growing grass. In treating for ants apply this product thoroughly to mounds. If possible, wet down treated area after applying insecticide to lawns. Keep children and pets off treated areas until insecticide is washed into the soil and the grass is thoroughly dry. Do not contaminate food or feed crops.**CITRUS** (Grapefruit, Kumquats, lemons, limes, oranges, tangerines and tangeloes): To control ants in citrus groves apply 15 - 20 pounds of this product per acre as a broadcast soil treatment when no fruit is present. To control termites apply 1 ounce of this product per tree to area to be banked and an additional 1-ounce to mound when bank is half completed.**BEANS:** To control Ants, Cutworms, Mole crickets, and White-fringed beetle grubs use at the rate of 20 lbs. per acre. Do not apply to green or snap beans after pods begin to form.**EGGPLANTS:** To control Ants, Cutworms, Mole Crickets, and White-fringed beetle grubs use at the rate of 20 lbs. per acre. Apply as a preplanting soil application or at time of planting only.**BROCCOLI:** To control Ants, Cutworms, Mole crickets, and White-fringed beetle grubs use at the rate of 20 lbs. per acre. Apply as a preplanting soil application or at time of planting only.**LETTUCE:** To control Ants, Cutworms, Mole Crickets and White-fringed beetle grubs use at the rate of 20 lbs. per acre. Apply as a preplanting soil application or at time of planting only.**CABBAGE:** To control Ants, Cutworms, Mole crickets, and White-fringed beetle grubs use at the rate of 20 lbs. per acre. Apply as a preplanting soil application or at time of planting only.**PEPPERS:** To control Ants, Cutworms, Mole crickets and White fringed beetle grubs use at the rate of 20 lbs. per acre. Apply as a preplanting soil application or at time of planting only.**COLLARDS:** To control Ants, Cutworms, Mole Crickets and White-fringed beetle grubs use at the rate of 20 lbs. per acre. Apply as a preplanting soil application or at time of planting only.**STRAWBERRIES:** To control Ants, Cutworms, Mole Crickets and White-fringed beetle grubs use at the rate of 20 lbs. per acre. Do not apply after fruit begins to form.**CUCUMBERS:** To control Wireworms and White grubs use at the rate of 25 lbs. per acre. Apply as a preplanting soil application or at time of planting only. May be injurious to some varieties.**SWEET CORN:** To control wireworms and White grubs use at the rate of 25 lbs. per acre. Apply as a preplant soil application or at time of planting only.

