

All dosages given on this label are for dilute applications only. Spray often enough to keep new growth covered. Use the higher dosage during periods of frequent and heavy rainfall.

APPLES, PEARS: To control powdery mildew and scab use 6 to 8 lbs. per 100 gals. of water in the cover sprays. Reduce dosage to 2 lbs. per 100 gals. of water if hot weather threatens. A micron sized sulphur is preferred in the pre-bloom through first cover sprays.

PEACH: To control brown rot and scab use 6 to 8 lbs. per 100 gals. water in pink, bloom, shuck, cover and pre-harvest sprays.

PLUM, control brown rot and leaf spot use 6 to 8 lbs. per 100 gals. shuck, cover and pre-harvest sprays.

own rot and leaf spot use 6 to 8 lbs. per 100 gals. water fall, shuck, cover and post-harvest sprays.

to control rust mite use 3 to 10 lbs. per 100 gals. of water. at first sign of damage and repeat as necessary.

To control powdery mildew and rust use 6 to 8 lbs. per acre in sufficient water for thorough coverage. Repeat applications in 7 to 14 day intervals as necessary.

NOTICE: When using lead arsenate with Sulphur Smoke Wettable always add an equal amount of Spray Lime, or other safener, as would be used when applying the Arsenical without sulphur. When using Zinc Sulfate and Spray Lime as a safener on peaches, always add Zinc Sulfate to the spray tank first, followed by Spray Lime, Lead Arsenate and Sulphur Smoke Wettable in order given.

July 24/16
124-001

NOTICE: Follow directions carefully. Time and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors, in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

W. R. GRACE & CO.
AGRICULTURAL PRODUCTS DIVISION
BALTIMORE, MARYLAND
NET WEIGHT POUNDS