

EPA REG. NO. 2124-68



TOTAL.

KEEP OUT OF REACH OF CHILDREN

This product is poisonous to fish and wildlife. Birds feeding on treated areas will be harmed. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on this label. Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

NOTE: Do not permit children or pets to go on the treated lawn until the insecticide has been washed into the soil by drenching or sprinkling with water and grass is completely dried.
Birds feeding on treated areas may be killed

NET WEIGHT 4 POUNDS

directions given herewith


UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

2124-681

Ornamenta Turf

Grain Sorghum (seed)
Corn (seed)
Wheat (seed)
Soybeans (seed)

(Seed)
Cotton (Seed)
Rice (Seed)



DIRECTIONS FOR USE (continued)

TERMITE CONTROL

To control termites in building: Mix with water at the rate of 1 pint Naco Chlordane 8 E C Insecticide to 12 gallons water and apply as follows:

1. Slab-Type Construction

- Apply 1 gallon per 10 sq. ft. as an overall pre-construction treatment under the slab and under attached porches.
- Apply 2 gallons per 5 lineal feet to critical areas under the slab as pre-construction treatment such as along the inside of foundation walls and around utility entrances and interior partition foundation walls.
- Apply 2 gallons per 5 lineal feet along the outside of the foundation using a trench to the depth of the foundation.

2. Crawl-Space Houses and Homes with Basements

- Apply 2 gallons per 5 lineal feet as a trenching application to critical areas under the house, such as along the inside of foundation walls and around piers and utility entrances. If trench is deeper than 15-18 inches, apply 4 gallons per 5 lineal feet.
- Apply 2 gallons per 5 lineal feet in a trench along the outside of foundation walls, including entrance platforms and porches where the foundation is not more than 15-18 inches deep. If over 15-18 inches deep but not over 30 inches deep, apply 4 gallons per 5 lineal feet of trench.
- Apply 1 gallon per 10 square feet of soil surface as a pre-construction treatment where the attached porches, entrance platforms and utility entrances have covering slab on fill of ground.

NOTE: For buildings with deep footings, dig trench 30 inches deep. Do not dig trench below level of footing.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.



CHLORI

ACTIVE INGREDIENTS:

Technical Chlordane*
Xylene

INERT INGREDIENTS

* Equivalent to 43.76% Octachloro-4, 7-methanot

KEEP OUT

HARMFUL IF SWALLOWED. AVOID BREATH SYMPTOMS. In case of contact with skin, Foodstuffs.

ANTIDOTE: In case of spillage on skin wash prometic of 1 tablespoon mustard in tumbler of wa This product is poisonous to fish and wildlife. Bi and streams. Do not apply in any manner not sp of spray from crop areas treated. Rinse equipmer away from water supplies. Containers should be

W.
AGR

Amarillo, Texas • Baltimore, Md. • Charleston, S
Lansing, Mich.

NET C





USDA Reg. No. 2124-742

CHLORDANE 8 E C INSECTICIDE



INGREDIENTS:

Technical Chlordane*	72.93%
Xylene	21.07%
Other Ingredients	6.00%
TOTAL	100.00%

*Equivalent to 43.76% Octachloro-4, 7-methanotetrahydroindane and 29.17% Related Compounds.

WARNING!

KEEP OUT OF REACH OF CHILDREN

IF SWALLOWED. AVOID BREATHING SPRAY MIST. CONTACT WITH SKIN CAN CAUSE TOXIC EFFECTS. In case of contact with skin, wash with soap and water. Avoid Contamination of Feed and Water.

NOTE: In case of spillage on skin wash promptly with soap and water, in case material has been swallowed — give 1 tablespoon mustard in tumbler of water. Call a Physician.

This product is poisonous to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds, streams. Do not apply in any manner not specified on this label. Do not apply when weather conditions favor drift from crop areas treated. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

W. R. GRACE & CO.
AGRICULTURAL CHEMICALS
MEMPHIS, TENNESSEE

El Paso, Texas • Baltimore, Md. • Charleston, S.C. • Columbus, Ohio • Ft. Pierce, Fla. • Henry, Ill. • Joplin, Mo. •
Lansing, Mich. • Nashville, Tenn. • New Albany, Ind.

NET CONTENTS GALLONS



GENERAL DIRECTIONS

Naco Chlordane 8 E C Insecticide is a concentrated insecticide and must be diluted in sufficient water for application in accordance with your State Agricultural Experiment Station Extension Service.

During prolonged exposure in spray-mixing and loading operations, wear clothing and a mask or respirator of a type found suitable by the U. S. Department of Health against Chlordane. Use this material only for recommended purposes and purposes. Hydraulic sprayers or low volume ground applicators may be used to apply.

DIRECTIONS FOR USE

PEPPERS: To control wireworms, apply 10 pints (in West) 4-8 pints (in East) and thoroughly work into upper 6-9 inches. Do not apply after planting time.

POTATO: To control white-fringed beetle grubs apply 5 pints per acre broadcast and thoroughly work into upper 3 inches. Do not feed tubers from Chlordane treated soil to livestock. Do not apply after planting.

To control white grubs apply 4-10 pints broadcast on soil and thoroughly work into upper 3-4 inches. Do not feed tubers from Chlordane treated soil to livestock. Do not apply after planting.

To control wireworms (except Southern potato wireworm in South where it is common) apply 4-8 pints per acre broadcast on soil and thoroughly work into upper 6-9 inches. Do not feed tubers from Chlordane treated soil to livestock. Do not apply after planting.

SWEET CORN: To control rootworms apply 1 pint per acre to row or hill at planting and immediately work into upper 3-4 inches. Do not apply after planting.

TOMATO, OUTDOOR: To control wireworms (in East) apply 4-8 pints per acre broadcast and thoroughly work into upper 4-6 inches. Do not apply after planting.

TURNIP: To control wireworms (in East) apply 4-8 pints per acre broadcast and thoroughly work into upper 4-6 inches. Do not apply after planting.

WATERMELON: To control wireworms apply 10 pints (in West) — apply 4-8 pints (in East) broadcast on soil and thoroughly work into upper 6-9 inches. Do not apply after planting.

