


Science



Multi-Purpose

# GARDEN DUST

or Spray

for **TOMATOES** & other **VEGETABLES**

**FRUITS - FLOWERS - SHRUBS - TREES**

**ACTIVE INGREDIENTS:**

	BY WEIGHT
Methoxychlor Technical <sup>1</sup>	5.00%
Malathion <sup>2</sup>	5.00%
Captaf <sup>3</sup>	4.89%
Related Derivatives	5.01%
<b>INERT INGREDIENTS:</b>	50.00%
<b>TOTAL</b>	<b>100.00%</b>

<sup>1</sup>Equivalent to 4.4%, 2,2 bis (p-methoxyphenyl)-1,1,1-trichloroethane and 0.6% other isomers and related products.

<sup>2</sup>O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

<sup>3</sup>N-(trichloromethylthio)-4-cyclohexene-1, 2 dicarboximide.

EPA Reg. No. 2125-13

**KEEP OUT OF REACH OF CHILDREN**  
**DANGER PELIGRO**

**PRECAUCION AL USUARIO:** Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

**STATEMENT OF PRACTICAL TREATMENT**

**IF IN EYES:** Flush with plenty of water. Call a physician.

**IF SWALLOWED:** Drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available drink large quantities of water. Avoid alcohol.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

**IF INHALED:** Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

**IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and water. Launder clothing before reusing.

**SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.**

Science Products Division  
MILLER CHEMICAL & FERTILIZER CORPORATION  
Hagerstown, Pennsylvania 17331

NET WEIGHT: 12 OUNCES

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS: DANGER**—Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Avoid contact with skin and clothing. Keep children and pets out of treated areas until spray has dried or dust has settled. Do not contaminate food or feed. Wash hands thoroughly after handling. Food utensils, such as tablespoons, should not be used for food after use with pesticides.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

**PHYSICAL OR CHEMICAL HAZARDS**

Avoid contact with open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Wear goggles or face shield when handling. Remove contaminated clothing and separately launder clothing before reuse.

Wear chemical resistant gloves when using indoors or outdoors. When using wear long pants and long-sleeved shirt and apply with the wind to your back. Wash non-disposable gloves thoroughly with soap and water before removing.

**STORAGE & DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal. Store in tightly closed original container in a locked area away from children and domestic animals. Do not reuse empty container. Securely wrap closed container in several layers of newspaper and discard in trash.

**DIRECTIONS**

Begin applications when insects or diseases first appear. Repeat as needed, usually at 5 to 10 day intervals. Apply thoroughly to all plant surfaces.

**DUSTING** - Apply in early morning, or late afternoon or evening, when air is still.

**SPRAYING** - Mix 10 tablespoons (this Dust with a little water to make a thin, smooth paste. Add paste to water to make 1 gallon; mix thoroughly. Agitate spray mix before and during spraying to maintain good dispersion.

**NOTE:** Do not apply to any food crop within 7 days of harvest unless otherwise noted. Remove residues remaining at harvest on edible portions of food crops by washing, wiping, brushing or other means.

**VEGETABLES**

**TOMATOES** - Controls aphids, flea beetles, leafhoppers, fall armyworms, blister beetles, tomato fruitworms, spider mites. Also anthracnose, early & late blight, septoria leaf spot. Begin applications after plants have been set into the field.

**CANTALOPES, CUCUMBERS** - Controls aphids, cucumber beetles, flea beetles, leafhoppers, spider mites; also angular leaf spot, anthracnose, downy mildew. Apply only when plants are dry.

**FLOWERS, SHRUBS, SHADE TREES**

Controls aphids, bagworms, birch leaf miner, boxwood leaf miner, lace bugs, leafhoppers, rose chafers, tarnished plant bugs, tent caterpillars, thrips, red spider mites. To control black spot on roses apply often enough to maintain coverage of new growth.

**FRUITS and BERRIES**

Apply at petal fall and every 10 to 14 days thereafter until control is achieved, unless otherwise directed. **NOTE:** Do not use within 10 days of an oil spray. Apply only as a spray. Consult your state Agricultural Extension Service for exact timing.

**APPLES, PEARS:** Make first application when leaves begin to appear. Repeat at 7 to 14 day intervals up to within 7 days of harvest except during bloom. Controls aphids, bud moths, codling moth, Forbes scale, Japanese beetles, pear psyllid, plum curculio, red-banded leafroller, spider mites; also Apple scab, bitter rot, black rot, Brooks spot.

**PEACHES:** Make first application when buds begin to show pink. Apply again when petals fall. Repeat at 7 to 10 day intervals until 21 days before harvest. Controls aphids, European red mites, Oriental fruit moth, plum curculio, red spider mites; also brown rot and peach scab.

**CHERRIES, PLUMS:** Apply as for peaches above. Controls aphids, cherry maggot, cherry fruit-fly, fruit-tree leafroller. Also controls brown rot and leaf spot on cherries.

**STRAWBERRIES, GRAPES:** Make first application in pre-bloom stage before infestation appears. Repeat as needed, usually at 10 to 14 day intervals; more frequently if needed. Controls aphids, grape berry moth, flea beetles, Japanese beetles, leafhoppers, mealybugs, spider mites, spittlebugs, strawberry weevil, strawberry leafroller; also botrytis fruit rot. Do not apply to grapes within 14 days of harvest nor on strawberries within 3 days of harvest.

**LIMITED WARRANTY:** Seller makes no warranty or guarantee of any kind, express or implied, concerning effects of use of this product, other than those specified on this label. Buyer, or User, accepts all responsibility for results due to misuse of this product.

77M MP

Prod. No. 82102

**BEST AVAILABLE COPY**

1/1 2125-13 PA 12

121