

READ ENTIRE LABEL CAREFULLY BEFORE USE.  
OBSERVE ALL CAUTIONS. USE ONLY AS DIRECTED.

#### DIRECTIONS FOR USE

**MIXING.** Science FRUIT-TREE and BERRY SPRAY is a wettable powder. Mix thoroughly with water as follows:

4 tablespoons (1 oz.) in gallon  
8 oz. in 8 gallons  
1 lb. in 16 gallons

**SPRAYING.** Start applications before insects become numerous and before disease symptoms appear. Repeat applications as needed to maintain effective control, usually at 7 to 14 day intervals, except as noted. Spray all exposed surfaces of plant thoroughly. Keep mixture agitated while spraying to maintain effective dispersion of particles.

#### FRUITS AND BERRIES

Do not use within 14 days of an oil spray. Consult state Agricultural Experiment Station or state Extension Service for exact timing.

**APPLES, PEARS.** Make first application when leaves begin to appear. Repeat at 7 to 14 day intervals up to within 7 days of harvest, except during bloom. Controls Aphids, Bud Moth, Codling Moth, Forbes Scale, Japanese Beetles, Pear Psylla, Plum Curculio, Red-banded Leafroller, Red Spider Mites; also Apple Scab, Bitter Rot, Black Rot, Brooks Spot.

**PEACHES.** Make first application when buds begin to show pink. Apply again when petals fall. Repeat at 7 to 10 day intervals until 21 days before harvest. Controls Aphids, European Red Mites, Oriental Fruit Moth, Plum Curculio, Red Spider Mites; also Brown Rot and Peach Scab.

**CHERRIES, PLUMS.** Apply as for PEACHES above. Controls Aphids, Cherry Maggot, Cherry Fruit-fly, Fruit-tree Leafroller. Also controls Brown Rot, Leaf Spot on cherries. Applications may be made up until 7 days before harvest.

**STRAWBERRIES, GRAPES.** Make first application in pre-bloom stage, before infestation first appears. Repeat as needed, usually at 10 to 14 day intervals; more frequently if needed. Do not apply to grapes within 14 days of harvest, nor on strawberries within 3 days of harvest. Controls Aphids, Grape Berry Moth, Flea Beetles, Japanese Beetles, Leafhoppers, Mealybugs, Spider Mites, Spittlebugs, Strawberry Weevil, Strawberry Leafroller; also Botrytis Fruit Rot.

#### HIGHLY EFFECTIVE

This Methoxychlor-Malathion-Captan combination is a highly effective, one-formula control for many common diseases, pests of certain fruits, berries, vegetables, flowers, and ornamentals.

#### VEGETABLES

**TOMATOES.** Controls Aphids, Flea Beetles, Leafhoppers, Fall Armyworm, Blister Beetles, Tomato Fruitworm. Also Early Blight, Late Blight, Septoria Leaf Spot, Anthracnose. Start applications after plants have been set into field. Do not apply within 7 days of harvest. Wash fruit thoroughly before eating.

**BEANS.** Controls Aphids, Mexican Bean Beetles. Do not apply after pods begin to form.

**CUCUMBERS, MELONS, SQUASH.** Controls Aphids, Cucumber Beetles, Flea Beetles, Leafhoppers, Spider Mites; also Angular Leaf Spot, Anthracnose, Downy Mildew. Apply only when plants are dry. Do not apply within 7 days of harvest.

**PEAS.** Controls Aphids. Do not apply within 7 days of harvest.

**NOTE.** Remove residues remaining at harvest from edible portions of fruits and vegetables by thoroughly washing, wiping, brushing or other effective means.

#### FLOWERS, SHRUBS, SHADE TREES

Controls Aphids, Bagworms, Birch Leaf Miner, Boxwood Leaf Miner, Lace Bugs, Leafhoppers, Rose Chafers, Tarnished Plant Bugs, Tent Caterpillars, Thrips, Red Spider Mites. For control of Black Spot on Roses apply often enough to maintain coverage of new growth.

**CAUTION!** Harmful if swallowed. Avoid breathing of dust or spray mist. Avoid repeated or prolonged contact with skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. This product is toxic to fish. Keep out of any body of water.

**NOTICE.** Seller makes no warranty of any kind, express or implied, concerning effects of use of this product other than those specified on this label. Buyer, or User, accepts all responsibility for results due to misuse of this product.

USDA Reg. No. 2125-14

Manufactured in U.S.A. by

SCIENCE PRODUCTS COMPANY, Inc.  
Chicago, Illinois

Science

# FRUIT-TREE AND BERRY SPRAY

NET WEIGHT  
1 LB.

Makes  
16 GALLONS



CONTROLS MANY DISEASES AND  
INSECTS OF FRUITS AND BERRIES

may also be used on  
**FLOWERS-SHRUBS-TREES  
VEGETABLE CROPS**

ACTIVE INGREDIENTS	
Technical Methoxychlor*	19.75%
Malathion**	7.50%
Captan***	15.00%
INERT INGREDIENTS	57.75%
TOTAL 100.00%	
*Equivalent to 17.40% of 2,2-bis [p-methoxyphenyl]-1,1,1-trichloroethane and 2.35% other isomers and related products.	
**O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate.	
***N-trichloromethylthio-tetrahydrophthalimide.	

2125-14  
2/9/1972  
Page 122

PA-PA  
307

ACCEPTED

FEB 9 1972

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. .... SUBJECT  
TO ATTACHED COMMENTS.

2125-14

2  
P