

2/25-20

6/20/1973

3-12-73

Michigan

Page 131

READ ENTIRE LABEL CAREFULLY BEFORE USE. OBSERVE ALL CAUTIONS. USE ONLY AS DIRECTED ON THIS LABEL.

FRUITS. Apples (7), apricots (21), blueberries (14), blackberries (14), cherries (7), grapes (14), peaches (21), pears (7), plums (7), gooseberries (14), cranberries (14), raspberries (14), strawberries (14). May injure McIntosh and Cortland apples if applied within four weeks of harvest.

Controls: Aphids, budmoth, cherry fruitfly, cherry fruitworm, codling moth, grape berry moth, grape leaf skeletonizer, Japanese beetle, Oriental fruit moth, red spider, plum curculio, spittlebug, strawberry weevil, San Jose scale (crawlers), leafhoppers, pear psylla.

IMPORTANT RESTRICTIONS — Observe carefully. Do not spray food crops within the number of days of harvest shown in parentheses for each crop listed. Edible foods should be washed, wiped, or brushed at harvest to remove possible residues. Do not apply to cucurbits unless plants are dry.

CAUTION! HARMFUL IF SWALLOWED. Avoid breathing of spray mist. Avoid contact with skin; wash thoroughly after skin contact and after using. Avoid contamination of feed, foodstuffs, cooking utensils.

This product is toxic to fish. Keep out of any body of water. This product is toxic to bees. Do not apply when bees are active. Do not re-use empty container. Destroy by breaking and burying with waste.

NOTICE. Seller makes no warranty of any kind, express or implied, concerning effects of use of this product other than those specified on this label. Buyer, or User, accepts all responsibility for results due to misuse of this product.

ACCEPT

JUN 20 1973
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 14320 SUBJECT TO ATTACHED COMMENTS.

Manufactured in U.S.A. by
SCIENCE PRODUCTS COMPANY
Chicago, Illinois

Science GARDEN INSECT Spray

KILLS GARDEN INSECTS!

as listed on this label

FLIES, MOSQUITOES, too!

ACTIVE INGREDIENTS	
*Malathion	12.5%
**Methoxychlor technical	12.5%
Xylene	69.0%
INERT INGREDIENTS	
Total	100.0%

7methyl dithiophosphate of diethyl mercaptosuccinate
Equivalent to 11.0% bis (p-methoxyphenyl)-1, 1, 1-trichloro-
ethane and 1.5% other isomers and related compounds.

Net Contents: 1 pint

READ ENTIRE LABEL CAREFULLY BEFORE USE. OBSERVE ALL CAUTIONS. USE ONLY AS DIRECTED ON THIS LABEL.

DIRECTIONS FOR USE

Add 2 tablespoons (1 oz.) per gallon of water or 1½ teaspoons per quart. Apply spray thoroughly to both upper and lower plant surfaces as soon as insects appear. Start spraying fruit trees at petal fall. Repeat as needed at 7 to 14 day intervals. For scale insects, use double strength. Do not spray food crops within the number of days of harvest shown in parentheses for each crop listed.

FLOWERS — ORNAMENTALS — TREES. Roses, azaleas, begonias, camellias, chrysanthemums, dahlias, gladioli, zinnias, evergreens, shade trees.

Controls: Aphids, tent caterpillars, Japanese beetle, red spider mites, leafhoppers, leafminers, lacebugs, mealybugs, thrips, tarnished plant bug, whitefly, bagworms, scale insects (crawlers). May cause injury to ferns, maidenhair, peris and crassula.

FLIES, MOSQUITOES—Outdoors only. Also ants, clover mites, crickets, earwigs, roaches and spiders around the home and in refuse piles. Use 6 tablespoons (3 oz.) in enough water to make 1 quart; or 1 quart in 2 gallons and 3 quarts water. Spray thoroughly surfaces on which insects alight. Repeat as needed.

VEGETABLES. Tomatoes (7), beans (3), cabbage (7), cauliflower (7), carrots (14), eggplant (7), peas (7), lettuce (14), peppers (7), potatoes (7), cucumbers (7), melons (7), squash (7), spinach (14).

Controls: Aphids, bean leaf beetle, blister beetle, imported cabbage worm, Colorado potato beetle, fall armyworm, flea beetles, Japanese beetle, cucumber beetle, leafhopper, Mexican bean beetle, tomato fruitworm, webworms, spider mites, lacebug, thrips.

EPA Reg. No. 2125-20-AA

CAUTION! KEEP OUT OF REACH OF CHILDREN
See also caution on back panel