

Flowers, Trees, Ornamentals. To control aphids, spider mites, lacebugs, white fly, scales (crawler stage), Japanese beetles, bagworms, mealybugs: use 1½ to 2 teaspoons per gallon (1½-2 pints per 100 gallons). Do not spray sensitive ferns such as Boston, Maidenhair, Pteris, and certain species of Crassula.

FLY CONTROL. Outdoors only. Around yards and homes. Use 5 tablespoons per gallon (1 pint in 6 gallons) and spray to run-off point on surfaces where flies alight: walls, fences, garbage pails, refuse piles, etc. Do not use inside dwellings.

MOSQUITOES, FLIES, Small Flying Insects. Outdoors. Use 4½ oz. (9 tablespoons) per gallon of water, fuel oil, or diesel oil and spray yard, lawn, patio or other outdoor area using a fine mist spray. Repeat as needed. Do not apply oil solutions to valuable ornamentals; injury may occur.

CAUTION. Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin; wash thoroughly after skin contact and after using. Avoid contamination of feed, foodstuffs, cooking utensils. Do not re-use empty container. Destroy by burying with waste or burning. Stay away from smoke or fumes.

This product is toxic to fish. Keep out of any body of water.

NOTICE. Seller makes no warranty of any kind, express or implied, concerning effects of use of this product other than those specified on this label. Buyer, or User, accepts all responsibility for results due to misuse of this product.

Science

57%

MALATHION

Spray

**Kills: Aphids, Mites, Mosquitoes
FLIES and GARDEN INSECTS**
as listed under Directions

**On: FLOWERS, VEGETABLES, FRUITS,
SHADE TREES, EVERGREENS, LAWNS**

ACTIVE INGREDIENTS

*Malathion 57%
Xylene 34%

INERT INGREDIENTS 9%
Total 100%

*O, O-dimethyl dithiophosphate
of diethyl mercaptosuccinate

Net Contents: 1 Pint

**READ ENTIRE LABEL CAREFULLY BEFORE USE.
OBSERVE ALL CAUTIONS. USE ONLY AS DIRECTED.
DIRECTIONS FOR USE**

Mix with water as directed. Spray upper and lower plant surfaces thoroughly when insects are first observed. Repeat applications as needed. Do not spray food crops within the number of days of harvest shown in parentheses for each crop listed.

Vegetables. To control aphids, spider mites on beans (1), broccoli (3), brussels sprouts (7), cabbage (7), cauliflower (7), celery (7), cucumbers (1), eggplant (3), kale (7), kohlrabi (7), lettuce (14), melons (1), potatoes (0), spinach (7), squash (1), tomatoes (1) and pea aphids, Mexican bean beetle, onion thrips on peas (3), beans (1) and onions (3) use 2 teaspoons per gallon (1½ pints per 100 gallons). Do not apply to cucurbits unless plants are dry.

Fruits. On apples (3) to control European red mite, green apple aphid, codling moth, plum curculio, red-banded leafroller use 2 to 3 teaspoons per gallon (1½-2 pints per 100 gallons). May cause injury to McIntosh and Cortland varieties if applied within 4 weeks of harvest. On pears (1) to control mites, pear psylla, mealybug use 2-3 teaspoons per gallon (1-2 pints per 100 gallons). May cause injury to Bosc pears. To control aphids, mites on strawberries (3); leafhoppers, mites, mealybugs on grapes (3) use 2 teaspoons per gallon (1½ pints per 100 gallons).

Manufactured in U.S.A. by

SCIENCE PRODUCTS COMPANY, Inc.
Chicago, Illinois

EPA Reg. No. 2125-22-AA

June 1972
With Comments

ACCEPTED WITH COMMENTS