

Science

KELTHANE

EC

Kills **RED SPIDER, MITES** on
ROSES, EVERGREENS, STRAWBERRIES
Flowers, Vegetables, Fruits, Berries, Nuts,
Shade Trees, Ornamentals, Nursery Stock

Kills CLOVER MITES in homes

CAUTION: KEEP OUT OF REACH OF CHILDREN

See other cautions on back panels

ACCEPTED

Science

KELTHANE

EC

RED SPIDER, MITES

Flowers, Vegetables, Fruits, Berries, Nuts
Shade Trees, Ornamentals, Nursery Stock

Kills **CLOVER MITES** in homes

CAUTION: KEEP OUT OF REACH OF CHILDREN

See other cautions on back panels

SCIENCE PRODUCTS COMPANY, Inc. 2640 No. Greenview Ave. Chicago, Illinois 60614

READ ENTIRE LABEL CAREFULLY BEFORE USE.
OBSERVE ALL CAUTIONS. USE ONLY AS DIRECTED.
DIRECTIONS FOR USE.

Use 2 teaspoons Kelthane EC in 1 gallon water. When mites first appear spray thoroughly both upper and lower plant surfaces. Repeat as necessary.

FLOWERS, ORNAMENTALS, NURSERY STOCK, TREES
Controls two-spotted mite (red spider), European red mite, Spruce mite, and other mites common on both greenhouse plants and outdoor plants such as roses, evergreens, shade trees, flowers, nursery stock, ornamentals. **Note:** Do not use on sensitive varieties or on tender foliage which may not tolerate emulsion sprays. Winter foliage may be more sensitive than summer foliage.

CLOVER MITES In Homes. Spray walls, foundations, windowsills; also plants and lawn at base of infested building.
VEGETABLES. Controls two-spotted mite (red spider), Pacific, Tropical and Tomato russet mites on beans, cucumbers, cantaloupes, melons, watermelons, pumpkins, winter and summer squash, tomatoes, ~~peppers~~ peppers. Do not use on beans within 7 days of harvest, nor on other crops in this section within 2 days of harvest. ←

RAWBERRIES. Controls two-spotted mite (red spider), strawberry mite (Atlantic mite), and six-spotted mite. For cyclamen mite use 2 to 4 teaspoons per gallon of water and drench plants thoroughly at 10 to 20 day intervals in spring when new growth starts. Or use as spot treatment during season. Do not use within 2 days of harvest.

NOTICE. Seller makes no warranty of any kind, express or implied, concerning effects of use of this product other than those specified on this label. Buyer, or User, accepts all responsibility for results due to misuse of this product.

← USDA REG. NO. 2125-25

BERRIES, GRAPES, NUTS. Controls Brown Almond (clover), two-spotted (red spider), European red, Pacific, McDaniel, Willamette, and yellow mites on grapes, raspberries, dewberries, loganberries, boysenberries, ~~walnuts~~ walnuts, filberts, pecans, chestnuts, hickory nuts. Apply just before bloom or whenever mites appear. ~~On nuts do not apply within 14 days of harvest.~~ On ~~nuts~~ nuts do not apply within 14 days of harvest. Do not use on grapes within 7 days of harvest, nor on berries within 2 days of harvest.

FRUITS. APPLES, PEARS, CRABAPPLES, QUINCES. For European red mite, clover mite, two-spotted (red spider), Schoene, McDaniel, yellow (carpini), Willamette, Pacific, and apple rust mites. Apply in second or later cover sprays or whenever mites appear. Do not apply within 7 days of harvest. **PEACHES, APRICOTS, NECTARINES, CHERRIES, PLUMS, PRUNES, FIGS.** For European red, clover (brown almond), two-spotted (red spider), Schoene, McDaniel, Pacific, Willamette, peach silver, plum, nursery, and yellow mites. Apply at petal fall to first cover spray or whenever mites appear. If repeat application is necessary, wait 30 days to avoid excessive residues. Do not apply to cherries, plums, prunes, figs within 7 days of harvest, nor on

2-2-72

2125-25
Comments

Controls two-spotted mite (red spider), European red mite, Spruce mite, and other mites common on both greenhouse plants and outdoor plants such as roses, evergreens, shade trees, flowers, nursery stock, ornamentals. Note: Do not use on sensitive varieties or on tender foliage which may not tolerate emulsion sprays. Winter foliage may be more sensitive than summer foliage.

CLOVER MITES In Homes. Spray walls, foundations, windowsills; also plants and lawn at base of infested building. VEGETABLES. Controls two-spotted mite (red spider), Pacific, Tropical and Tomato russet mites on beans, cucumbers, cantaloupes, melons, watermelons, pumpkins, winter and summer squash, tomatoes, ~~peppers~~ peppers. Do not use on beans within 7 days of harvest, nor on other crops in this section within 2 days of harvest. ←

RAWBERRIES. Controls two-spotted mite (red spider), strawberry mite (Atlantic mite), and six-spotted mite. For cyclamen mite use 2 to 4 teaspoons per gallon of water and drench plants thoroughly at 10 to 20 day intervals in spring when new growth starts. Or use as spot treatment during season. Do not use within 2 days of harvest.

NOTICE. Seller makes no warranty of any kind, express or implied, concerning effects of use of this product other than those specified on this label. Buyer, or User, accepts all responsibility for results due to misuse of this product.

USDA REG. NO. 2125-25 ←

BERRIES, GRAPES, NUTS. Controls Brown Almond (clover), two-spotted (red spider), European red, Pacific, McDaniel, Willamette, and yellow mites on grapes, raspberries, dewberries, loganberries, boysenberries, ~~walnuts~~ walnuts, filberts, pecans, chestnuts, hickory nuts. Apply just before bloom or whenever mites appear. ~~On nuts do not apply within 14~~

On ~~nuts~~ nuts do not apply within 14 days of harvest. Do not use on grapes within 7 days of harvest, nor on berries within 2 days of harvest.

FRUITS. APPLES, PEARS, CRABAPPLES, QUINCES. For European red mite, clover mite, two-spotted (red spider), Schoene, McDaniel, yellow (carpini), Willamette, Pacific, and apple rust mites. Apply in second or later cover sprays or whenever mites appear. Do not apply within 7 days of harvest. PEACHES, APRICOTS, NECTARINES, CHERRIES, PLUMS, PRUNES, FIGS. For European red, clover (brown almond), two-spotted (red spider), Schoene, McDaniel, Pacific, Willamette, peach silver, plum, nursery, and yellow mites. Apply at petal fall to first cover spray or whenever mites appear. If repeat application is necessary, wait 30 days to avoid excessive residues. Do not apply to cherries, plums, prunes, figs within 7 days of harvest; nor on other fruits listed above within 14 days of harvest. CITRUS FRUITS. For citrus red (purple), citrus flat, six-spotted and Yuma mites, on non-bearing trees or as post-harvest application on bearing trees, follow timing recommendations obtainable from state agricultural experiment station.

CAUTION. Harmful if swallowed. Avoid: inhalation of spray mist; skin contact; contamination of feed, foodstuffs. Wash thoroughly after handling. In case of skin contact, wash with soap and water.

This product is toxic to fish. Keep out of any body of water.

Distributed by

SCIENCE PRODUCTS COMPANY, Inc.
Chicago, Illinois

2-2-72

2125-25
Comments