

SCIENCE 1% ROTENONE

DUST OR SPRAY FOR  
VEGETABLES, BERBERIES, FLOWERS

MAY BE USED RIGHT UP TO DAY BEFORE HARVEST!

ACTIVE INGREDIENTS:

|                         |        |
|-------------------------|--------|
| Rotenone-----           | 1.0%   |
| Other Cube Resins-----  | 1.6%   |
| INERT INGREDIENTS:----- | 97.4%  |
| TOTAL-----              | 100.0% |

EPA Reg. No. 2125-26

EPA Est. No. 2125

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Call a physician or Poison Control Center. Do not induce vomiting or give anything by mouth to an unconscious person. IF INHALED: Remove from exposure. IF IN EYES: Flush with plenty of water. Get medical attention. IF ON SKIN: Remove contaminated clothing. Wash affected area with soap and water. Launder clothes before reusing.

See Side Panel For Additional Precautionary Statements.

NET WEIGHT: 1 POUND, 4 POUNDS

SCIENCE PRODUCTS DIVISION  
MILLER CHEMICAL & FERTILIZER CORPORATION  
Hanover, Pennsylvania 17331

BEST DOCUMENT AVAILABLE

2125 210

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS-CAUTION: Harmful if swallowed. May cause eye, skin, nose and throat irritation. Avoid contact with skin, eyes and clothing. Avoid inhalation of dust. Wash thoroughly after handling. Food utensils such as teaspoons and cups should not be used for food after use with pesticides. ENVIRONMENTAL HAZARD: This product is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of waste. Do not apply directly to water or wetlands.

GENERAL INFORMATION

This product kills weed seeds in the ground as they germinate, before they emerge from the soil. It does not kill weeds after they emerge. For full effectiveness, before applying this product the soil should be thoroughly cultivated to remove existing weeds.

CONTROLS: crabgrass, other annual grasses, and certain broad-leaved weeds as follows: annual bluegrass, barnyard grass, browntop panicum, carpetweed, common chickweed, dodder, fall panicum, Florida pussley, goosegrass, green foxtail, hairy crabgrass, Johnsongrass (from seed), lambsquarters, lovegrass, Palmer amaranth, purslane, prostrate spurge, red-root pigweed, spotted spurge, stinkgrass, Texas millet, witchgrass, Wright groundcherry, yellow foxtail.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Store in the tightly closed, original container in a locked storage area away from children and pets. Do not reuse empty container. Securely wrap closed container in several layers of newspaper and discard in trash.

NOT DOCUMENT AVAILABLE

Apply as soon as insects appear. Repeat applications at 5 to 10 day intervals. Be sure to cover thoroughly both upper and lower surfaces of foliage as well as stems. Rotenone may be used until the day before harvest.

**AS A DUST** Apply directly from this container through the sifter top or apply with a dust applicator.

**AS A SPRAY** Make a paste by stirring 4 heaping tablespoons in a little water. Mix this paste with water to make a gallon of spray. When spraying, shake sprayer occasionally to maintain an effective suspension.

**VEGETABLES**

Asparagus, beans, leafy table, broccol, brussels sprouts, cabbage, carrots, cauliflower, collard, cucumbers, eggplant, kale, kohlrabi, lettuce, melons, mint, mustard greens, okra, peppers, pumpkins, radishes, rutabagas, squash, swiss chard, tomatoes, turnips.

**Controls:** Aphids (except waxy cabbage aphid), asparagus beetle, twelve spotted, bean leaf beetles, imported cabbage worm, Colorado potato beetles, cucumber beetles, flea beetle, Mexican bean beetle, pepper maggots, squash vine borer, tobacco weevils.

**BERRIES and SMALL FRUITS**

Blackberry, blueberry, boysenberry, huckleberry, raspberry, strawberry, gooseberry, grape, Loganberry, Marionberry, Rubus.

**Controls:** Blueberry maggot, raspberry beetle, gooseberry fruitworm, imported European spruce sawfly, Japanese beetles, raspberry cane borers, raspberry cane worms, raspberry sawfly, and necked cane borers, strawberry leaf beetles.

**FLOWERS and ORNAMENTALS**

Use as dust or spray on flowers, ornamentals, shrubs, etc. Control certain aphids, Japanese Beetles, etc.

ORIGINAL WARRANTY: 1 YEAR  
PRINTS NUMBER: 123456789

BEST DOCUMENT AVAILABLE