


Reg # 2125-26  
PM 1-1  
1-1/2



# 1% ROTENONE

DUST OR SPRAY FOR  
VEGETABLES, BERRIES, FLOWERS  
MAY BE USED RIGHT UP TO  
DAY BEFORE HARVEST!

ACTIVE INGREDIENTS:	
Rotenone .....	1.0%
Other Cube Resins .....	1.6%
INERT INGREDIENTS: .....	97.4%
TOTAL .....	100.0%

EPA Reg. No. 2125-26

EPA Est. No. 904-NJ-2

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**

### STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Call a physician or Poison Control Center. Do not induce vomiting or give anything by mouth to an unconscious person. IF INHALED: Remove from exposure. IF IN EYES: Flush with plenty of water. Get medical attention. IF ON SKIN: Remove contaminated clothing. Wash affected area with soap and water. Launder clothes before reusing.

SEE BACK PANEL FOR ADDITIONAL  
PRECAUTIONARY STATEMENTS.

SCIENCE PRODUCTS DIVISION  
MILLER CHEMICAL & FERTILIZER CORPORATION  
Hanover, Pennsylvania 17331

NET WEIGHT 1 POUND

DEC 3 1981

*Handwritten signatures and notes*

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS:**

**CAUTION**—Harmful if swallowed. May cause eye, skin, nose and throat irritation. Avoid contact with skin, eyes and clothing. Avoid inhalation of dust. Wash thoroughly after handling. Food utensils such as teaspoons and cups should not be used for food after use with pesticides. **ENVIRONMENTAL HAZARD:** This product is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of waste. Do not apply directly to water or wetlands.

**DIRECTIONS FOR USE:**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. Store in the tightly closed, original container in a locked storage area away from children and pets. Do not reuse empty container. Securely wrap closed container in several layers of newspaper and discard in trash.

**APPLICATION DIRECTIONS**

Apply as soon as insects appear. Repeat applications at 5 to 10 day intervals. Be sure to cover thoroughly both upper and lower surfaces of foliage as well as stems. Rotenone may be used until the day before harvest.

**AS A DUST:** Apply directly from this container through the sifter top; or apply with a dust applicator.

**AS A SPRAY:** Make a paste by stirring 4 heaping tablespoons in a little water. Mix this paste with water to make a gallon of spray. When spraying, shake sprayer occasionally to maintain an effective suspension. **DO NOT APPLY TO FRUITS OR VEGETABLES ON THE DAY OF HARVEST.**

**VEGETABLES:**

Asparagus, beans, beets (table), broccoli, brussels sprouts, cabbage, carrots, cauliflower, collard, cucumbers, eggplant, kale, kohlrabi, lettuce, melons, mint, mustard greens, okra, peppers, pumpkins, radishes, rutabagas, squash, swiss chard, tomatoes, turnips.

**Controls:** Aphids (except waxy cabbage aphid), asparagus beetles (twelve-spotted), bean leaf beetles, imported cabbage worms, Colorado potato beetles, cucumber beetles, flea beetles, Mexican bean beetles, pepper maggots, squash vine borers, vegetable weevils.

**BERRIES AND SMALL FRUITS:**

Blackberry, blueberry, boysenberry, cherry, currant, dewberry, gooseberry, grape, loganberry, raspberry, strawberry.

**Controls:** Blueberry maggots, cherry fruitflies, flea beetles, gooseberry fruitworms, imported currant worms, Japanese beetles, raspberry can borers, raspberry fruitworms, raspberry sawfly, red-necked cane borers, spittle bugs, strawberry leaf beetles.

**FLOWERS, ORNAMENTALS, SHRUBS, VINES:**

Use as dust or spray on flowers, ornamentals, shrubs, or vines to control certain aphids, Japanese Beetles, lace bugs, rose slugs.

**LIMITED WARRANTY:** BUYER ASSUMES ALL RISKS OF USE, STORAGE AND HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HERewith.