

DIRECTIONS FOR USE

Mix 4 teaspoons per gallon of water or 4 oz. in 20 gallons unless otherwise specified. Apply with ordinary sprayer or hose-connected sprayer. Spray all plant surfaces thoroughly.

ROSES, OTHER FLOWERS

ROSES. Black Spot, Powdery Mildew. Apply as soon as new leaves first appear. Repeat as needed at 7-10 day intervals during the season.

CARNATIONS. Alternaria Leaf Spot, Rust. Begin 2 weeks after planting. Make 3 to 4 applications two weeks apart.

CHRYSANTHEMUMS, IRIS. Septoria Leafspot, Didymellina Leaf Spot. Spray immediately after planting. Repeat at 7-10 day intervals, as needed, to maintain effective control.

MARIGOLDS, ZINNIAS, CHRYSANTHEMUMS, ASTERS, PHLOX, SNAPDRAGONS. Alternaria Leafspot, Anthracnose, Powdery Mildew. Spray weekly during periods of rapid new growth and severe infection. Spray twice weekly if severe infection is due to high humidity and frequent rains.

TOMATOES, OTHER VEGETABLES

TOMATOES. Anthracnose. Use 8 teaspoons per gallon. Apply when blossoms are open. Repeat at 7-10 day intervals until fruits are harvested.

MELONS, SQUASH, PUMPKINS. Anthracnose, Downy Mildew, Powdery Mildew. Apply when first true leaves appear. Repeat weekly until fruits are harvested.

FRUITS, BERRIES

APPLES. Alternaria Leafspot, Bitter Rot, Black Rot, Brooks Fruit Spot, Fly Speck, Scab, Scaly Blotch. Begin applications at delayed-

EPA Reg. No. 2125-27-AA

dormant period. Spray thoroughly at 7-14 day intervals until harvest. In western states do not apply between bloom and July 1 to avoid injury. Russetting may be caused on sensitive varieties in Eastern states when used just before bloom to 30 days after petal fall.

BLACKBERRIES, RASPBERRIES. Fruit Rot, Gray Mold, Spur Blight. Apply before first infection. Repeat at 7-day intervals until harvest.

STRAWBERRIES. Leaf Spot, Fruit Rot. Start spraying before blooms appear. Repeat at 7-day intervals until berries are harvested.

COMPATIBILITY. May be used in combination with most commonly used insecticides, fungicides, and adjuvants except strongly alkaline materials such as hydrated lime.

CAUTION. Avoid inhalation of dust or spray mist. Avoid contact with skin. Avoid contamination of feed or foodstuffs. Do not use in combination with, or soon after using, oil sprays. Do not reuse container. Destroy when empty.

This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as directed on this label.

NOTICE. Buyer, or User, assumes all risks of use, storage, or handling of this material not in strict accordance with directions given on this label.

ACTIVE INGREDIENT

*FOLPET (N-trichloromethylthio-phthalimide) 75%
INERT INGREDIENTS 25%

Total 100%

Distributed by
SCIENCE PRODUCTS COMPANY, Inc.
Chicago, Illinois



Rose and GARDEN FUNGICIDE

for
BLACK SPOT, POWDERY MILDEW
and certain other diseases

on
**ROSES, OTHER FLOWERS, FRUITS,
VEGETABLES, TOMATOES**

contains **75% PHALTAN**
Phthalimide Fungicide

PHALTAN is a registered trademark for phthalimide fungicide
J.C. RICHMOND
CHICAGO, ILL. 60602

CAUTION: KEEP OUT OF REACH OF CHILDREN
See other cautions on back panel

ACCEPTED
May 1, 1973

Mix 4 teaspoons per gallon of water or 4 oz in 20 gallons unless otherwise specified. Apply with ordinary sprayer or hose-connected sprayer. Spray all plant surfaces thoroughly.

ROSES. Black Spot, Powdery Mildew. Apply as soon as new leaves first appear. Repeat as needed at 7-10 day intervals during the season.

CARNATIONS. Alternaria Leaf Spot, Rust. Begin 2 weeks after planting. Make 3 to 4 applications two weeks apart.

CHRYSANTHEMUMS, IRIS. Septoria Leafspot, Didymellina Leaf Spot. Spray immediately after planting. Repeat at 7-10 day intervals, as needed, to maintain effective control.

MARIGOLDS, ZINNIAS, CHRYSANTHEMUMS, ASTERS, PHLOX, SNAPDRAGONS. Alternaria Leafspot, Anthracnose, Powdery Mildew. Spray weekly during periods of rapid new growth and severe infection. Spray twice weekly if severe infection is due to high humidity and frequent rains.

TOMATOES. Anthracnose. Use 8 teaspoons per gallon. Apply when blossoms are open. Repeat at 7-10 day intervals until fruits are harvested.

MELONS, SQUASH, PUMPKINS. Anthracnose, Downy Mildew, Powdery Mildew. Apply when first true leaves appear. Repeat weekly until fruits are harvested.

APPLES. Alternaria Leafspot, Bitter Rot, Black Rot, Brooks Fruit Spot, Fly Speck, Scab, Scoty Blotch. Begin applications at delayed

EPA Reg. No. 2125-27-AA

dormant period. Spray thoroughly at 7-14 day intervals until harvest. In western states do not apply between bloom and July 1 to avoid injury. Russetting may be caused on sensitive varieties in Eastern states when used just before bloom to 30 days after petal fall.

BLACKBERRIES, RASPBERRIES. Fruit Rot, Gray Mold, Spur Blight. Apply before first infection. Repeat at 7-day intervals until harvest.

STRAWBERRIES. Leaf Spot, Fruit Rot. Start spraying before blooms appear. Repeat at 7-day intervals until berries are harvested.

May be used in combination with most commonly used insecticides, fungicides, and adjuvants except strongly alkaline materials such as hydrated lime.

Avoid inhalation of dust or spray mist. Avoid contact with skin. Avoid contamination of feed or foodstuffs. Do not use in combination with, or soon after using, oil sprays. Do not reuse container. Destroy when empty.

This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as directed on this label.

Buyer, or User, assumes all risks of use, storage, or handling of this material not in strict accordance with directions given on this label.

ACTIVE INGREDIENT

*FOLPET (N-trichloromethylthio-phthalimide) 75%

INERT INGREDIENTS 25%

Total 100%

Distributed by
SCIENCE PRODUCTS COMPANY, Inc.
Chicago, Illinois



**Rose and
GARDEN
FUNGICIDE**

for
BLACK SPOT, POWDERY MILDEW
and certain other diseases

on
**ROSES, OTHER FLOWERS, FRUITS,
VEGETABLES, TOMATOES**

contains **75% PHALTAN ***
Phthalimide Fungicide

Net Weight: 8 oz.

*PHALTAN is a registered trade mark for phthalimide fungicide.

U.S. Pat. Nos
2,553,770-2,553,771

CAUTION: KEEP OUT OF REACH OF CHILDREN
See other cautions on back panels

Handwritten: Mar 1, 1973