edge, 6 in. wide, to reduce growth the following Spring. Apply to green, growing grass just before it becomes dormant. This grass will green up about 2 weeks later in Spring because of the growth inhibition effect of the chemical.

NOTE. Color of grass is not usually affected at recommended application rates. Treated grass may be greener than untreated grass for a few weeks; it may also remain greener during drought.

CAUTION. May be harmful if swallowed. Avoid inhalation of spray mist. Avoid contact with skin, eyes, clothing. Do not use on food or feed crops. Do not use within 12 hours of expected rain. Use only on grass established 3 or more years. Not for general lawn application. Use only for edging.

NOTICE: Seller makes no warranty of any kind, express or implied, concerning effects of use of this product other than those specified on this label. Buyer, or User, accepts all responsibility for results due to misuse of this product.

SCIENCE PRODUCTS COMPANY, Inc.
Chicago, Illinois



STOP-GRASS

The Chemical Grass Edger

SAVES WORK OF MOWING GRASS along lawn edges, driveways, walks

KEEPS GRASS SHORT!

Grass Needs Little or No Trimming

Treats 8,000 Linear Feet, 6 inches Wide

Part Carret I port

NO MOST CHARGE CONTY COMPANIES. DESPRE IF MY COMPANIES, PORTECNING OR COMMEN MA. DOMY OR CHICAGO IN A MAST PLACE.

FOR USE ON GRASS EDGES

Grass stays short; grows so slowly it needs no trimming for several months.

Where to use. Use on grass lawn edges along flower-beds, driveways, hedges, fences; around ornaments, trees, shrubs; on waste areas to keep down grass height. Use special care in spraying to minimize drift on flowers, trees, shrubs.

FOLLOW THESE DIRECTIONS CAREFULLY SPRING APPLICATION (April, May): Mix 2 table-spoons (1 oz.) in 1 gal. water. Apply with hand sprayer or sprinkling can to 500 ft. of lawn edge, 6 inches wide. Hose-end sprayers: set applicato to apply 1 oz. to 250 sq. ft. (500 ft., 6 in. wide).

Compressed air sprayers: Use 4 tablespoons in 1 gal. water and apply uniformly to 800 ft., 6 inches wide. Uniform application to full area being treated is very important.

Best time to apply: The 2-week period when grass has grown 2-4 inches. At this time forsythia and dandelions are in full bloom. If later application is made, it may be desirable to make a single mowing one week after spray is applied.

FALL APPLICATION (Oct. 1- Nov. 15). Apply 4 tablespoons (2 oz.) in 1 gal. water to 500 ft. of lawn

USDA Rea. No. 2125-31

edge, 6 in. wide, to reduce growth the following Spring. Apply to green, growing grass just before it becomes dormant. This grass will green up about 2 weeks later in Spring because of the growth inhibition effect of the chemical.

NOTE. Color of grass is not usually affected at recommended application rates. Treated grass may be greener than untreated grass for a few weeks; it may also remain greener during drought.

CAUTION. May be harmful if swallowed. Avoid inhalation of spray mist. Avoid contact with skin, eyes, clothing. Do not use on food or feed crops. Do not use within 12 hours of expected rain. Use only on grass established 3 or more years. Not for general lawn application. Use only for edging.

NOTICE: Seller makes no warranty of any kind, express or implied, concerning effects of use of this product other than those specified on this label. Buyer, or User, accepts all responsibility for results due to misuse of this product.

ACTIVE INGREDIENT

Diethanolamine salt of 1,2-INERT INGREDIENTS42%

*Equivalent to 30% Maleic Hydrazide

*U.S. Patent No. 2,575,954 Distributed by



The Chemical Grass Edger

SAVES WORK OF MOWING GRASS along lawn edges, driveways, walks

KEEPS GRASS SHORT!

Grass Needs Little or No Trimming

Treats 8,000 Linear Feet, 64nches Wide

Net Contents: 1 pint

OF HISE ON OPAS SHOPE

Grass stays short; grows so slowly it needs no trimming for several months.

Where to use. Use on grass lawn edges along flower-beds, driveways, hedges, fences; around ornaments, trees, shrubs; on waste areas to keep down grass height. Use special care in spraying to minimize drift on flowers, trees, shrubs.

FOLLOW THESE DIRECTIONS CAREFULLY SPRING APPLICATION (April, May): Mix 2 tablespoons (1 oz.) in 1 gal. water. Apply with hand sprayer or sprinkling can to 500 ft. of lawn edge, 6 inches wide. Hose-end sprayers: set applicato to apply 1 oz. to 250 sq. ft. (500 ft., 6 in. wide).

Compressed air sprayers: Use 4 tablespoons in 1 gal. water and apply uniformly to 800 ft., 6 inches wide. Uniform application to full area being treated is very important.

Best time to apply: The 2-week period when grass has grown 2-4 inches. At this time forsythia and dandelions are in full bloom. If later application is made, it may be desirable to make a single mowing one week after spray is applied.

FALL APPLICATION (Oct. 1- Nov. 15). Apply 4 tablespoons (2 oz.) in 1 gal. water to 500 ft. of lawn

