




BERRY-SET

A HORMONE-TYPE SPRAY FOR **BIGGER BERRIES**



**JUST
SPRAY
THE
PLANTS**

**BIGGER
LUSCIOUS
LONGER
BEARING**

STRAWBERRIES

CAUTION: KEEP OUT OF REACH OF CHILDREN



**UP TO 57%
LARGER BERRIES**



Manufactured in U.S.A. by
SCIENCE PRODUCTS COMPANY, Inc.
Chicago, Illinois

DIRECTIONS FOR USE

Dissolve two level teaspoons ($\frac{1}{4}$ oz.) BERRY-SET powder in one gallon water, or 1 ounce in 4 gallons. Spray with this solution using ordinary, clean spray equipment. Hand, knapsack, or power sprayer may be used.

When To Apply. Make first application about 14 to 16 days after full bloom. At this time most of the flowers are open and there will be many small fruits (about $\frac{1}{4}$ to $\frac{1}{3}$ full size) on the plants from the first flowers. Blossoms and fruits should be thoroughly wetted by the spray mist. Early morning or late evening applications are best. Make a second application in 4 or 5 days. Make no application within 14 days of harvest.

Large Scale Application. When using agricultural-type power sprayers at 50 to 100 lbs. pressure, apply 100 gallons per acre. Knapsack-type sprayers will require considerably less solution per acre.

Net Weight: 3 oz.

ACTIVE INGREDIENT	
Beta-naphthoxyacetic acid	2.67%
INERT INGREDIENTS	97.33%
	TOTAL 100.00%

USDA Reg. No. 2125-39

NOTICE. Conditions of use being beyond seller's control, seller makes no warranty or guarantee of any kind, express or implied, concerning effects of use of this product. Buyer assumes all risks of use whether in accordance with directions or not.

Jan. 23, 1968