

Read entire label thoroughly before use. Use strictly according to label directions.
Observe **CAUTION** and **LIMITATION** statements carefully.

DIRECTIONS — CROPS

Apply this dust when insects or the damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surfaces of all foliage. Repeat as necessary, usually at 7 day intervals. Do not apply in excess. Observe preharvest limitations.

HOME GARDENS Apply with dust gun, or from shaker top can.

ACREAGE APPLICATIONS. Apply **40 lbs. per acre** to all crops listed on this label except as follows: Cucumbers, melons, pumpkins, squash (**20 lbs. per acre**), Peas (**30 lbs. per acre**); Beans, green lima, and snap (**42 1/2 lbs. per acre**), Grapes, peppers (**60 lbs. per acre**), Eggplant, tomatoes (**80 lbs. per acre**).

VEGETABLE INSECTS

ASPARAGUS Controls asparagus beetle, Apache cicada. Apply during cutting season. Do not repeat application within 3 days.

BEANS Green, lima, and snap beans, cowpeas, black-eyed peas. Controls Mexican bean beetle, bean leaf beetle, cucumber beetles, flea beetles, Japanese beetles, leafhoppers, velvet bean caterpillar, western bean cutworm, armyworms, corn earworm, lima bean pod borer, stink bugs, tarnished plant bugs.

SWEET CORN Controls corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetles, Japanese beetle, sap beetles and leafhoppers. For larvae or whorl (bait) worm damage, and foliage feeders, apply to entire plant. Repeat as necessary.

For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are essential for effective control.

CUCUMBER, MELONS, PUMPKIN, SQUASH Controls pickleworm, melonworm, cucumber beetles, flea beetles, leafhoppers, squash bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or during prolonged high humidity. SEVIN injures watermelons in Florida.

TOMATO, POTATO, EGGPLANT, PEAS, PEPPERS. For Colorado potato beetle, flea beetles, leafhoppers, European corn borer, fall armyworm, lace bugs, tomato fruit worm, tomato hornworm, tarnished plant bugs, stink bugs.

OKRA. For corn earworm, stink bugs.

CARROTS, CHINESE CABBAGE, COLLARDS, DANDELIONS, ENDIVE (ESCAROLE), GARDEN BEETS, HORSERADISH, KALE, LETTUCE, MUSTARD GREENS, PARSLEY, PARSNIPS, RADISHES, RUTABAGAS, SALSIFY, SPINACH, SWISS CHARD, TURNIPS. For flea beetles, harlequin bug, leafhoppers, armyworms, cabbage caterpillars, corn earworm, spittlebugs, tarnished plant bugs, stink bugs. To control small cabbage loopers, use on a 5 to 7 day schedule.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI. For flea beetles, harlequin bug, armyworms, cabbage caterpillars, corn earworm. To control small cabbage loopers, use on a 5 to 7 day schedule.

SMALL FRUIT CROPS

GRAPES For grape berry moth, Japanese beetles, June beetles, red banded leafroller, grape leaf folder, grape leafhopper, grape leaf skeletonizer. Apply just before first brood leaf folder larvae emerges from rolls and as needed for leafhoppers.

STRAWBERRIES For meadow spittlebug, strawberry leaf folder, strawberry weevils.

BLUEBERRIES For blueberry fruit borer, cherry and cranberry fruitworm.

BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, RASPBERRIES For Japanese beetles, leaf rollers, leafhoppers, raspberry aphids.

GRASSHOPPERS ON ABOVE CROPS Apply as often as necessary on these crops to control grasshoppers.

FLOWERS — SHRUBS — TREES

This dust can safely be used on a wide variety of flowers, shrubs, and small trees, such as **ROSES, CARNATIONS, GLADIOLUS, CHRYSANTHEMUM, LILAC, ARBOR VITAE, JUNIPER, HYDRANGEA, AZALEA, ELM, MAPLE, OAK, DOGWOOD, BIRCH, PINES.** Do not use on **BOSTON IVY.**

For such pests as apple aphid, bagworm, birch leaf miner, blister beetles, boxelder bug, boxwood leaf miner, elm leaf aphid, elm leaf beetle, flea beetles, gypsy moth, Japanese beetle, June beetles, lace bugs, leafhoppers, leaf roller, mealy bugs, mimosa webworm, oak leaf miners, orange tortrix, periodical cicada, plant bugs, puss caterpillars, psyllids, rose aphid, rose slug, scale insects, tent caterpillars, thorn bugs, thrips (exposed), willow leaf beetles.

PETS

ON DOGS, CATS For ticks and fleas, dust liberally over the entire animal, avoiding the eyes, and rub thoroughly into the skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Avoid treatment of kittens less than 4 weeks old.

IN and AROUND DOG and CAT QUARTERS. For ticks and fleas, dust liberally in and around sleeping quarters, kennels, and dog houses by means of a shaker can, hand or dust gun.

LIMITATIONS OF USE BEFORE HARVEST

Allow the designated time limit shown between application and harvest of the crops listed below.

1 day: Asparagus, strawberries.

3 days: Broccoli, Brussels sprouts, cabbage, cauliflower, head lettuce, kohlrabi, garden beets (roots), horseradish, parsnips, radishes, rutabagas, salsify (roots), turnip (roots).

7 days: Blackberries, boysenberries, dewberries, loganberries, raspberries.

14 days: Chinese cabbage, collards, dandelion, endive (escarole), garden beets (tops), kale, leaf lettuce, mustard greens, parsley, salsify (tops), spinach, Swiss chard, turnips (tops).

CAUTION

HARMFUL IF SWALLOWED OR INHALED. Avoid prolonged or repeated breathing of dust. Do not take internally. Avoid prolonged or repeated contact with skin. Wash thoroughly after using. Avoid contamination of food. Toxic to honey bees, avoid use on plants during flowering and pollination. Injury to tender foliage may occur if applications are made when plants are wet or in the presence of high humidity of several days duration following treatment.

Toxic to Fish and Shellfish. Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

NOTICE. Seller makes no warranty of any kind, express or implied, concerning effects of use of this product other than those specified on this label. Buyer, or User, accepts all responsibility for results due to misuse of this product.

U.S.D.A. Reg. No. 2125-40

Manufactured in U.S.A. by
SCIENCE PRODUCTS COMPANY, Inc.
Chicago, Illinois

Apply before or after
harvest.

BEANS - For Colorado potato beetle, European corn borer, and other insects. Apply as often as necessary, usually 1 to 2 times per week. Observe preharvest interval.

CUCUMBERS - For cucumber beetle, leafhoppers, and other insects. Apply as often as necessary, usually 1 to 2 times per week. Observe preharvest interval.

PEPPERS - For Colorado potato beetle, European corn borer, and other insects. Apply as often as necessary, usually 1 to 2 times per week. Observe preharvest interval.

VEGETABLES - For Colorado potato beetle, European corn borer, and other insects. Apply as often as necessary, usually 1 to 2 times per week. Observe preharvest interval.

FRUIT TREES - For Colorado potato beetle, European corn borer, and other insects. Apply as often as necessary, usually 1 to 2 times per week. Observe preharvest interval.

FLOWERS - For Colorado potato beetle, European corn borer, and other insects. Apply as often as necessary, usually 1 to 2 times per week. Observe preharvest interval.

Apply before or after
harvest.

STRAWBERRIES - For meadow spittlebug, strawberry leaf miner, and strawberry weevil.

BLUEBERRIES - For blueberry fruit worm, and cranberry fruit worm.

BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, RASPBERRIES - For Japanese beetles, leaf rollers, leafhoppers, and raspberry aphids.

GRASSHOPPERS ON ABOVE CROPS - Apply as often as necessary on these crops to control grasshoppers.

FLOWERS — SHRUBS — TREES

This dust can safely be used on a wide variety of flowers, shrubs, and small trees, such as **ROSES, CARNATIONS, GLADIOLUS, CHRYSANTHEMUM, LILAC, ARBOR VITAE, JUNIPER, HYDRANGEA, AZALEA, ELM, MAPLE, OAK, DOGWOOD, BIRCH, PINES**. Do not use on **BOSTON IVY**.

For such pests as apple aphid, bagworm, birch leaf miner, blister beetle, boxelder bug, boxwood leaf miner, elm leaf aphid, elm leaf beetle, flea beetles, gypsy moth, Japanese beetle, June beetle, lace bugs, leafhoppers, leaf miner, mealy bugs, mimosa webworm, oak leaf miners, orange tortrix, periodical cicada, plant bugs, puss caterpillars, psyllids, rose aphid, rose slug, scale insects, tent caterpillars, thorn bugs, thrips (exposed), willow leaf beetle.

PET USE

ON DOGS - CATS - For ticks and fleas dust liberally over the entire animal, avoiding the eyes, and rub thoroughly into the skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Avoid treatment of kittens less than 4 weeks old.

IN and AROUND DOG and CAT QUARTERS - For ticks and fleas dust liberally in and around sleeping quarters, kennels, and dog houses by means of a shaker can, hand or dust gun.

LIMITATIONS OF USE BEFORE HARVEST

Allow the designated time limit shown between application and harvest of the crops listed below.

- 1 day - Asparagus, strawberries
- 3 days - Broccoli, Brussels sprouts, cabbage, cauliflower, head lettuce, kohlrabi, garden beets (roots), horseradish, parsnips, radishes, rutabagas, salsify (roots), turnip (roots)
- 7 days - Blackberries, boysenberries, dewberries, loganberries, raspberries
- 14 days - Chinese cabbage, collards, dandelion, endive (escarole), garden beets (tops), kale, leaf lettuce, mustard greens, parsley, salsify (tops), spinach, Swiss chard, turnips (tops).

CAUTION

HARMFUL IF SWALLOWED OR INHALED - Avoid prolonged or repeated breathing of dust. Do not take internally. Avoid prolonged or repeated contact with skin. Wash thoroughly after using. Avoid contamination of food. Toxic to honey bees, avoid use on plants during flowering and pollination. Injury to tender foliage may occur if applications are made when plants are wet or in the presence of high humidity of several days duration following treatment.

NOTE TO PHYSICIAN - Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

NOTICE - Seller makes no warranty of any kind, express or implied, concerning effects of use of this product other than those specified on this label. Buyer, or User, accepts all responsibility for results due to misuse of this product.

Science

5%

SEV
Dust

Controls Many
VEGETABLES •
Flowers • Shrubs

ACTIVE INGREDIENT
Carbaryl (1-naphthyl N-methylcarbamate)
INERT INGREDIENTS

Net Weight:

*Registered trademark of Union Carbide
(1-naphthyl N-methylcarbamate) U.S.

125-40

Manufactured in U.S.A. by
SCIENCE PRODUCTS COMPANY, Inc.
Chicago, Illinois

CAUTION: KEEP OUT
See other cautions