

CONTROLS MANY INSECTS ON

VEGETABLES, SMALL FRUITS,
FLOWERS, SHRUBS and ORNAMENTALS
also Fleas & Ticks on Cats and Dogs
as listed on label

ACTIVE INGREDIENT:

Carbaryl (1 naphthyl methylcarbamate)-----5.0%
INERT INGREDIENTS:-----35.0%
TOTAL-----100.0%

EPA Reg. No. 2125-40

EPA Est. No. 2125-IL-1

Registered trademark of Union Carbide Corporation for carbaryl insecticide.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panel For Additional Precautionary Statements.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Call physician or Poison Control Center. Do not induce vomiting or give anything by mouth to an unconscious person. IF INHALED: Remove victim to fresh air. IF IN EYES: Flush with plenty of water. IF ON SKIN: Remove contaminated clothing. Wash affected area with soap and water. Launder clothing before reuse. NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Intravenous injection of atropine is antidotal.

BEST DOCUMENT AVAILABLE

11 41

PREPARATORY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS--CAUTION: Harmful if inhaled or swallowed. Do not brush hair. Avoid contact with skin, eyes or clothing. Do not take internally. Wear long-sleeves. Change clothing daily. Wash hands and face before eating. Wash thoroughly after handling. Keep children and pets off treated area until spray has dried.

ENVIRONMENTAL HAZARDS: Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from treated area. Do not contaminate water by cleaning of equipment or disposal of waste. **BE CAUTION: MAY KILL BEEKEEPERS IN SUBSTANTIAL NUMBERS.** This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not use when value of bees as pollinators is more important than insect control. Warn beekeepers to locate hives beyond bee flight range until one week after application, or take equally effective precautions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Store in closed, original container in a locked area away from children and domestic animals. For disposal, securely wrap closed container in several layers of newspaper and place in trash collection. Do not reuse container.

USE OF SEVIN DUST

Apply this dust when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surfaces of all foliage. Repeat as necessary, usually at 7 day intervals. Do not apply in excess. Observe preharvest limitations.

HOME GARDENS. Apply with dust gun, or from shaker-top container. **ACREAGE APPLICATIONS:** Apply 60 lbs. per acre to all crops listed on this label except as follows: Cucumbers, melons, pumpkins, squash (20 lbs. per acre); Peas (30 lbs. per acre); Beans, green, lima, and snap (42 1/2 lbs. per acre); Grapes, peppers (60 lbs. per acre); Eggplant, tomatoes (80 lbs. per acre).

VEGETABLE INSECTS

ASPARAGUS. Controls asparagus beetle, Asparagus fly. Apply during cutting season. Do not repeat application within 3 days. **BEANS.** (Green, lima, and snap beans; cowpeas, black-eyed peas) Controls Mexican bean beetle, bean leaf beetle, cucumber beetle, flea beetle, Japanese beetle, leafhoppers, velvet-bean caterpillar, western bean cutworm, armyworms, corn earworm, lima bean pod borer, stink bugs, tarnished plant bug.

SWEET CORN. Controls corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetle, Japanese beetle, sap beetle and leafhoppers. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary.

For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are essential for effective control. Application to tassels during pollen shed will seriously reduce bee population.

CUCUMBER, MELONS, PUMPKIN, SQUASH. Controls pickleworm, melonworm, cucumber beetle, flea beetle, leafhoppers, squash bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or during prolonged high humidity. SEVIN inhibits watermelon in Florida.

TOMATO, POTATO, EGGPLANT, PEAS, PEPPERS. For Colorado potato beetle, flea beetle, leafhoppers, European corn borer, fall armyworm, lace bugs, tomato fruit worm, tomato horn worm, tarnished plant bug, stink bugs.

ORZA. For corn earworm, stink bug. **CABBAGE, CHINESE CABBAGE, COLLARDS, DANDIELIONS, EN-DIVE (SCAROLET), GARDEN BEETS, NORSEADISH, KALE, LETTUCE, MUSTARD GREENS, PARSLEY, PARSNIPS, RADISHES, RUTABAGAS, TARSIFY, SPINACH, SWISS CHARD, TURNIPS.** For flea beetle, harlequin bug, leafhoppers, armyworms, cabbage caterpillars, corn earworm, spittlebugs, tarnished plant bug, stink bugs. To control small cabbage leapers, use on a 5 to 7 day schedule.

CABBAGE BROCCOLI, BEANS, SPROUTS, CAULIFLOWER, KOMRABI. For flea beetle, harlequin bug, armyworms, cabbage caterpillars, corn earworm. To control small cabbage leapers use on a 5 to 7 day schedule.

SMALL FRUIT INSECTS

GRAPES. For grape berry moth, Japanese beetle, flea beetle, red-banded sphinx, grape leaf folder, grape leafhopper, grape leaf skeletonizer. Apply just before first brood leaf folder larvae emerges from cells and as needed for leafhoppers. **STRAWBERRIES.** For meadow spittlebug, strawberry leaf roller, strawberry weevil.

BLUESBERRIES. For blueberry maggot, cherry and cranberry fruitworm. **BLACKBERRIES, BOYSSONBERRIES, DEWBERRIES, LORANBERRIES, RASPBERRIES.** For Japanese beetle, leaf rollers, leafhoppers, raspberry aphids. **GRAPEHOPPERS ON ABOVE CROPS.** Apply as often as necessary on these crops to control grapehoppers.

FLOWERS — SHRUBS — TREES

This dust can safely be used on a wide variety of flowers, shrubs, and small trees, such as ROSES, CARNATIONS, GLADIOLUS, CHRYSANTHEMUM, LILAC, ABOR VITAE, RHODOS, HYDRANGEA, AZALEA, GUM, MAPLE, OAK, BIRCH, BURNING BUSH, PINES. Do not use on GORSEBERRY IVY.

For such pests as apple aphid, bagworm, birch leaf miner, blister beetle, boxelder bug, browned leaf miner, elm leaf aphid, elm leaf beetle, flea beetle, gypsy moth, Japanese beetle, June beetle, leaf beetle, leafhoppers, leaf roller, mealy bug, minute webworm, oak leaf miner, orange tortrix, periodical cicada, plant bug, rose caterpillar, psyllid, rose aphid, rose slug, scale insects, tent caterpillar, thorn bug, thrip (open), willow leaf beetle.

PET USE

ON DOGS, CATS. For ticks and flea dust liberally over the entire animal, avoiding the eyes, and rub thoroughly into the skin. Comb out dead fleas a few hours after treatment. Repeat at weekly intervals if needed. Avoid treatment of kittens less than 4 weeks old.

IN and AROUND DOG and CAT QUARTERS. For ticks and flea dust liberally in and around sleeping quarters, kennels, and dog houses by means of a shaker can, hand or dust gun.

LIMITATIONS OF USE BEFORE HARVEST

Allow the designated time (last shown) between application and harvest of the crops listed below:

1 day: Asparagus, strawberries. 3 days: Broccoli, Brussels sprouts, cabbage, cauliflower, head lettuce, kohlrabi, garden bean (root), horseradish, parsnip, radishes, rutabaga, salsify (root), turnip (root).

7 days: Blackberries, boysenberries, dewberries, loganberries, raspberries. 14 days: Chinese cabbage, collards, dandelion, endive (scarole), garden bean (leaf), kale, leaf lettuce, mustard greens, parsley, salsify (leaf), spinach, Swiss chard, turnip (leaf).

LIMITED WARRANTY: SEVIN AND ALL OTHERS ON THE MARKET ARE MADE BY THE NATIONAL CHEMICAL COMPANY, INC. THE NATIONAL CHEMICAL COMPANY, INC.