

Science FORE LAWN FUNGICIDE

**Controls Fungus Diseases
on lawns and certain ornamentals**

ACTIVE INGREDIENT

A co-ordination product of zinc ion and
manganese ethylene bisdithiocarbamate.....80%
in which the ingredients are:

Manganese + +16%

Zinc + + 2%

Ethylene bisdithiocarbamate
ion (C₄H₆N₂S₄) =62%

INERT INGREDIENT20%

Total.....100%

Net Weight 8 oz.

Treats up to 2,000 sq. ft. of lawn

®FORE, Registered Trade Mark, Rohm & Haas Company

Read Entire Label Thoroughly. Use Strictly According to Label Directions.

GENERAL INFORMATION. FORE is a new type fungicide which controls most common lawn grass diseases and major fungus diseases of certain ornamentals such as Roses, Glads, and Mums.

Best results are obtained by using a regular, protective spray program.

FORE is compatible with most commonly used pesticides. When used in combination with other pesticides, or nutritional sprays, the other materials should be added to the mix last.

DIRECTIONS FOR USE

LAWN and TURF DISEASES

Dollar Spot (Sclerotinia), Red Thread, Fusarium Blight (F. Roseum), Slime Molds, Copper Spot. In a protective spray program, use 6 to 8 oz. in sufficient water (5 gallons or more) to cover 1,000 sq. ft. of lawn area. Apply when grass greens up in the Spring; repeat at 7 to 14 day intervals.

When conditions especially favor disease development, use 8 oz. per 1,000 sq. ft. Apply every 7 days until disease threat is past.

Helminthosporium Melting Out, Rhizoctonia Brown Patch, Rust. Use 4 oz. in sufficient water (5 gallons or more) to cover 1,000 sq. ft. Apply when grass greens up in Spring. For Helminthosporium and Rust, repeat every 7 to 14 days; for Rhizoctonia, repeat every 7 days.

Pythium. Use 8 oz. in sufficient water to cover 1,000 sq. ft. Start applications at first sign of disease and repeat at 5 day intervals (oftener if conditions especially favor disease development). Continue until disease threat is past. FUSARIUM SNOW MOLD

Algae. Use 6 oz. in enough water to cover 1,000 sq. ft. Apply when algae first appear. Repeat every 7 days until threat is past.

Fusarium Snow Mold. Use 6 to 8 oz. in enough water to cover 1,000 sq. ft. Make applications during the winter at 2 to 6 week intervals.

ORNAMENTAL DISEASES

Mix 1½ level tablespoons FORE in a gallon of water. Spray all plant surfaces thoroughly.

ROSES. For Black Spot. Begin spraying when plants are well leafed out. Apply weekly.

CHRYSANTHEMUM. For Botrytis Petal Spot. Apply twice weekly during the blooming season.

GLADIOLUS. For leaf and flower spot diseases caused by Curvularia and Botrytis. Start regular weekly applications before diseases appear. Increase to 2 or 3 applications per week after diseases appear and during wet weather.

MISTLETOE. For Mold control. Dip cut mistletoe sprigs in a solution containing 1½ level tablespoons FORE and 3 drops Science SPREADER-STICKER per gallon of water. When sprigs are thoroughly wet, remove and air dry.

CAUTION. May cause irritation of nose, throat, eyes, and skin. Do not breathe dust or spray mist. In case of skin contact, wash with water; for eyes get medical attention. Do not contaminate fish ponds. Fish may be harmed.

IMPORTANT. Keep away from fire and sparks. Store in cool, dry place. Do not allow to become wet or overheated in storage. This may cause chemical changes and may generate flammable vapors. Keep container closed when not in use.

NON-WARRANTY. Seller makes no warranty of any kind, express or implied, concerning effects of use of this product other than those specified on this label. Buyer, or User, accepts all responsibility for results due to misuse of this product.

Distributed by

CAUTION: KEEP OUT OF REACH OF CHILDREN
See other cautions on back panels

U S D A Reg. No 2125-43

SCIENCE PRODUCTS COMPANY, Inc.,
Chicago, Illinois

ACCEPTED WITH COMMENTS

2125-43
REGISTERED
8/1/52