


ACCEPTED

SEP 23 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under  
EPA Reg. No. 2125-43



**FORE®**

**Lawn & Ornamental Fungicide**  
CONTROLS: BLACK SPOT ON ROSES, BROWN PATCH ON TURF AND OTHER LISTED PLANT DISEASES

**ACTIVE INGREDIENT:**

A coordination product of zinc, iron, and manganese ethylene	
bis(dithiocarbamate) in which the ingredients are	80%
Manganese + +	16%
Zinc + +	2%
Ethylene bis(dithiocarbamate)	62%
Iron + +	20%
<b>TOTAL</b>	<b>100%</b>

FORE® is a trademark of Rohm and Haas Co. U.S. Patent No. 3,379,610  
EPA Reg. No. 2125-43 EPA Est. No. 2125-IL-1

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**IF OF PRACTICAL TREATMENT**

IF SWALLOWED: Give 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Call physician or Poison Control Center.  
IF IN EYES: Flush with plenty of water. Get medical attention. IF INHALED: Remove victim to fresh air. IF ON SKIN: Remove contaminated clothing. Wash affected area with soap and water. Launder clothes before reusing. Get medical attention if irritation continues.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

SCIENCE PRODUCTS DIVISION  
MILLER CHEMICAL & FERTILIZER CORPORATION  
Hanover, Pennsylvania 17331

NET WEIGHT 8 OUNCES

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS—CAUTION:** May cause irritation of nose, throat, eyes, and skin. Do not breathe dust or spray mist. Avoid contact with skin, eyes and clothes. Wear rubber gloves when using indoors. When using outdoors, wear long pants, long sleeve shirt, and rubber gloves. Outdoors, spray with the wind to your back. Wash nondisposable gloves thoroughly with soap and water before removing. Keep children and pets out of treated areas until dry. Do not contaminate food or feed. Any utensils used for measuring, such as teaspoons, tablespoons, cups, etc., should not be used to measure food products after measuring pesticides. **ENVIRONMENTAL HAZARD:** This product is toxic to fish. Do not apply directly to water or wetlands. Do not apply where runoff is likely to occur or when weather conditions favor drift from treated area. Do not contaminate water by direct application or cleaning of equipment. **PHYSICAL AND CHEMICAL HAZARD:** Keep away from fire and sparks. Do not allow to become wet or overheated in storage. This may bring on a chemical change which will impair the fungicidal effectiveness and generate flammable vapors.

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with this label.

**STORAGE AND DISPOSAL:**

Store this product in its closed, original container in a locked area away from children and domestic animals. Do not reuse empty container. For disposal, securely wrap original container in several layers of newspaper and discard in trash.

**LAWN AND TURF:** Apply the directed amount of fungicide mixed in a sufficient volume of water to cover 1,000 sq. ft. of lawn area (5 to 10 gals.). Repeat as indicated until the disease symptoms disappear or conditions favoring spread of the turf disease have abated. Start application when grass greens up in the Spring or as directed. **ALGAE:** 6 ozs.—Apply when algae appear. Repeat at 7 day intervals until threat is past. **COPPER SPOT, FUSARIUM BLIGHT (F. ROSEUM), RED THREAD SLIME MOLDS:** 4-6 ozs. at 7 to 14 day intervals until threat passes; or 6-8 ozs. at 7 days in conditions of high disease development. **DOLLAR SPOT:** 6-8 ozs. at 7 to 14 day intervals. Use higher rate and shortest interval during periods of high development. **FUSARIUM SNOW MOLD:** 6-8 ozs. during winter at 2 to 6 week intervals. **HELMINTHOSPORIUM MELTING OUT, RHIZOCTONIA BROWN PATCH, RUST:** 4 ozs. 7 to 14 day intervals (7 days for Melting Out). **PYTHIUM BLIGHT:** 8 ozs. at 5 day intervals when disease first appears.

**DO NOT GRAZE TREATED AREA OR FEED CLIPPINGS TO LIVESTOCK.**

**ORNAMENTALS**

Mix 1½ level tablespoons per gallon of water. Apply as a cover spray at 7 to 10 day intervals throughout the season starting when plants are well leaved out.

**CARNATION—**Leaf Spot; **DAHLIA—**Botrytis blight; **HOLLY—**Purple spot; **HOLLYHOCK—**Anthraxnose, Leaf Spot; **RUST; HONEYSUCKLE—**Herpobasidium blight; **IRIS—**Leaf spot; **LILY—**Botrytis blight; **ORNAMENTAL CRABAPPLE—**Cedar-apple, Rust, Scab; **PANSY—**Anthraxnose; **ROSES—**Black spot; **SNAPDRAGON—**Rust; **TULIP—**Botrytis blight (lily); **ZINNIA—**Leaf blight. **AZALEAS, CAMELLIA, RHODODENDRON—**Petal blight. Spray 2 to 3 times a week while flowers are opening. Direct spray into flowers and spread ground under bushes thoroughly. **CHRYSANTHEMUM—**Botrytis Petal Spot. Apply twice weekly during the bloom season. **FLOWERING DOGWOOD—**Anthraxnose. Apply when buds begin to open, when bracts have fallen, 4 weeks later again in late summer after flower buds for next season formed. **GLADIOLUS—**For leaf and flower spot diseases caused by Botrytis and Curvularia apply 2 to 3 gallons of dilute spray per 1,000 sq. ft. depending upon the size of the plants. Flower Spikes, reduce spray dilution to ¼ tablespoons per gallon of water. Weekly applications before diseases appear. Increase to 2 to 3 applications per week after diseases appear and during wet weather. **PACHYSANDRA—**Volvetella blight. Apply a solution containing 4 level tbs. per gal. as a drench 50 sq. ft. of bed area. Make 5 applications at 10 to 14 day intervals starting at first sign of disease. **PEONY—**Botrytis Phytophthora blight. Apply in early spring and early drenching soil around plants as well as the foliage. Destroy infected parts promptly.

**USE SPREADER-STICKER TO IMPROVE WETTING AND DISTRIBUTION**

**NOTE:** This product is compatible with most commonly used pesticides. When used in combination with other pesticides, nutritional sprays, or spreader stickers, mix the fungicide half of the water in the sprayer. Premix the other materials and add the mixture to the spray tank. Stir well as balance of water is added.

**LIMITED WARRANTY:** BUYER ASSUMES ALL RISKS OF STORAGE AND HANDLING OF THIS MATERIAL NOT USED IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

47M MP

Prod No. 8

©Copyright - 1983 - Miller Chemical & Fertilizer Corporation

BEST AVAILABLE COPY