

READ ENTIRE LABEL THOROUGHLY BEFORE USE.
OBSERVE ALL CAUTIONS. USE ONLY AS DIRECTED.

DIRECTIONS FOR USE

Controls diseases indicated on plants listed.

Unless otherwise directed use 1½ level tablespoons per gallon of water to make spray mix. Thoroughly spray both upper and under surfaces of foliage. Unless otherwise directed begin when disease first appears; repeat every 7 to 10 days as needed. Spray more frequently if infestation is severe. When a number in parentheses follows the plant name, do not spray that crop within the number of days of harvest shown.

VEGETABLES

TOMATOES (5) • Anthrachnose, early and late blight, gray leaf spot (stemphylium), septoria leaf spot. Begin application when tomatoes on first cluster are well-formed. Do not use on young plants in greenhouse.

BEANS (4), green and dry. Downy mildew, rust. Begin when plants are small. Repeat every 5 to 7 days.

BROCCOLI • Alternaria leaf spot, downy mildew. If applied within 3 days of harvest remove residues by washing or trimming.

BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI • Alternaria leaf spot, downy mildew.

CABBAGE (7) • Alternaria leaf spot, downy mildew. Begin 7 to 10 days after planting. Repeat every 3 to 5 days.

CARROTS • Leaf spot diseases. Start spraying when plants are 6 weeks old or disease threatens.

CELERY (14) • Early and late blight. Start spraying when plants are established in the garden. Remove residues at harvest by stripping, trimming, and washing.

CUCUMBERS (5), **SQUASH** (5) • Anthrachnose, downy mildew, pythium fruit rot. Begin when disease first appears or runners start to form.

CANTALOUPE (5), **WATERMELON** (5) • Anthrachnose, downy mildew, cercospora leaf spot, gummy stem blight. Begin when vines begin to run or disease first appears.

EGGPLANT, PEPPERS • Anthrachnose, fruit rot. Apply as fruit begins to form. Repeat weekly through harvest.

LETTUCE (10), **ENDIVE** (10) • Downy mildew. Remove residues at harvest from head lettuce by stripping and trimming outer leaves; and from leaf lettuce and endive by washing.

NION • Botrytis leaf blight, downy mildew, purple blotch.

PUMPKINS • Angular leaf spot, downy mildew. Use 3 table-
spoons per gallon. Spray thoroughly.

SPINACH (10) • Downy mildew, white rust. Remove residues at harvest by washing.

FLOWERS

Begin spraying when new growth starts or when leaves first unfold. Apply at weekly intervals during season.

CARNATIONS, PANSIES: anthracnose, leaf spot. **DAHLIAS, ZINNIAS:** botrytis, leaf spot. • **FLOWERING DOGWOOD:** an-
thrachnose; apply when buds begin to open and again when
bracts fall, and 4 weeks later, and again in late summer.

GLADIOLUS: botrytis, curvularia, Stemphylium leaf spot.

PEONY: botrytis and phytophthora blights; apply in early
spring and early fall to foliage and to soil around plants;
repeat at 7 to 10 day intervals during growing season.

ROSES: black spot. **SNAPDRAGON:** rust.

APPLES (30)

Black rot, bitter rot, brown rot, bullseye rot, fly speck, rust,
scab, sooty blotch, white rot. Begin application in the pre-
pink sprays, repeat in subsequent pre-bloom, bloom and
cover sprays in accordance with local state spray schedules.
Do not spray within 30 days of harvest.

CAUTION. May cause irritation of nose, throat, eyes and
skin. Do not breathe dust or spray mist. May be absorbed
through skin. In case of skin contact wash with water, for
eyes, wash thoroughly with water and get medical attention.
This product is toxic to fish. Keep out of any body of water.
IMPORTANT. Keep away from fire and sparks. Store in a
cool, dry place. Do not allow to become wet or overheated
in storage. This may cause chemical changes which will im-
pair the effectiveness of this product and may also generate
flammable vapors. Keep container closed when not in use.
Do not reuse this container. Destroy it when empty.

NOTICE. Buyer, or User, assumes all risks of use, storage or
handling of this material not in strict accordance with the
directions given on this label.

Distributed by

Science

TOMATOES

and on
other Vegetables, Fruit, Flowers

ACTIVE INGREDIENT

Maneb (Manganese ethylene

bisdithiocarbamate)80%

INERT INGREDIENTS20%

TOTAL 100%

Net Weight 6 oz.

ACCEPTED