

2125-53

5/27/1970

Page 122



50%

METHOXYCHLOR

WETTABLE POWDER

DDT SUBSTITUTE

(For certain applications)

Kills certain insects on Vegetables,
Fruits, Flowers, Trees

Cankerworm, flea beetle, inchworm, Japanese beetle, leafhopper, rose chafer and other insects listed under directions.

ACTIVE INGREDIENTS	
*Methoxychlor, technical	50.0%
INERT INGREDIENTS	50.0%
TOTAL	100.0%

*Equivalent to 44% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane & 5% other isomers & related products.

Net Contents: 8 oz.

READ ENTIRE LABEL CAREFULLY BEFORE USE.

OBSERVE ALL CAUTIONS. USE ONLY AS DIRECTED ON THIS LABEL.

DIRECTIONS FOR USE

These directions are for home garden use only, not for large scale, agricultural applications. Thoroughly mix 2 tablespoons this product in 1 gallon water. First make a smooth paste by mixing powder with small amount of water. Then mix with more water to make 1 gallon spray mix. Apply spray thoroughly to all surfaces of plant. Begin applications at first sign of infestation; on fruits, begin at petal fall. Repeat at 7 to 14 day intervals, as needed, to maintain effective control. Do not spray food crops within the number of days of harvest shown in parentheses for each crop.

VEGETABLES

On ASPARAGUS(7), BEANS(3), BEETS*(7), BROCCOLI(14), BRUSSELS SPROUTS(14), CABBAGE(3), CANTALOUPE(1), CARROTS*(7), CAULIFLOWER(7), COLLARDS(14), CORN(7), CUCUMBERS(1), EGGPLANT(1), KALE(14), KOHLRABI(1), LETTUCE(14), MELONS(7), PEAS(7), PEPPERS(1), POTATOES(0), PUMPKIN(1), RADISHES(7), RUTABAGAS(7), SPINACH(14), SQUASH(1), SUMMER SQUASH(1), TOMATOES(1), TURNIPS*(7). Controls asparagus beetle, bean leaf beetle, blister beetle, imported cabbage worm, Colorado potato beetle, cucumber beetle, fall armyworm, flea beetle, Japanese beetle, leafhopper, Mexican bean beetle, pea weevil, squash vine borer.

*Beets, carrots, turnips. If tops are to be used for food, do not spray within 14 days of harvest.

FRUIT TREES

On APPLES(7), APRICOTS(21), CHERRIES(7), NECTARINES(21), PEACHES(21), PEARS(7), PLUMS(7), PRUNES(7), QUINCES(7). Controls apple maggot, cankerworm, cherry fruit fly,

cherry fruitworm, codling moth, Japanese beetle, leafhopper, pear slug, plum curculio, tent caterpillar.

SMALL FRUITS AND BERRIES

On BLACKBERRIES(14), BLUEBERRIES(14), CRANBERRIES(14), CURRANTS(14), GOOSEBERRIES(14), GRAPES(14), LOGANBERRIES(14), RASPBERRIES(14), STRAWBERRIES(14). Controls cranberry fruitworm, cherry fruitworm, grape berry moth, grape leaf skeletonizer, Japanese beetle, leafhopper, rose chafer, San Jose scale (crawlers), spittlebug, strawberry weevil.

ORNAMENTALS: Flowers, Trees, Shrubs

Controls fall armyworm, blister beetle, cankerworm, cucumber beetle, flea beetle, gypsy moth, inchworm, Japanese beetle, leafhopper, leafroller, rose chafer, rose slug, sawfly, spittlebug, tent caterpillar, tussock moth, webworm.

CAUTION. HARMFUL IF SWALLOWED. Avoid skin contact and breathing of spray mist. Wash thoroughly in case of skin contact and after using. Avoid contamination of feed and food-stuffs.

This product is toxic to fish. Keep out of any body of water.

Also toxic to bees. Do not apply where bees are active.

NOTICE. Buyer, or User, assumes all risks of use, storage, and handling of this material not in strict accordance with the directions given on this label.

USDA Reg. No. 2125-53

Distributed by

SCIENCE PRODUCTS COMPANY, INC.
Chicago, Illinois

CAUTION: KEEP OUT OF REACH OF CHILDREN
See other cautions on back panel.

ACCEPTED
MAY 27 1970
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
ED UNDER NO. 2125-535 SUBJECT
TO ATTACHED COMMENTS.

2125-535