

TREES, SHRUBS, ORNAMENTAL PLANTINGS

Trees, established plants. Apply to properly prepared soil early in Spring, or at a later time. Late summer application controls fall-germinating weeds.

Newly-set-out plants. Apply immediately after setting out, whether in late fall or in early spring.

weed control ## with
Especially recommended for/these
Flowers, Trees, Shrubs, Ornamentals

| | | |
|-----------------|--------------------|-------------------|
| Ash | Rhododendron | Fernoxone |
| Ash | Rose | Fernoxone |
| Ash | Russian olive | Forget me not |
| Ash | Spiraea | Four O'clock |
| Berberis | Spruce | Foxglove |
| Borwick | Sycamore | Gaillardia |
| Birch | Tree peony | Geranium |
| Camellia | Tulip tree | Gladiolus |
| Chestnut | Variiegated privet | Gold dust |
| Coltsfoot | Viburnum | Gold marguerite |
| Collinsia | Walden | Golden yucca |
| Cornflower | Weigela | Impatiens |
| Cypress | Willow | Lavender |
| Deutzia | Yew | Lily |
| Dogwood | Bulbifer | Lupine |
| Eleagnus | English ivy | Mangold |
| Elm | English ivy | Morning glory |
| Euonymus | Honeysuckle | Mother of thyme |
| Filix | Parthenocidia | Mourning bride |
| Forsythia | Wisteria | Mourning pink |
| Gum | Aperatum | Nasturtium |
| Hawthorn | Alyssum | Ornith |
| Holly | Aster | Peonies |
| Hydrangea | Baby's breath | Petunia |
| Jac. andromeda | Beilflower | Pink heart |
| Juniper | Bleeding heart | Pink plant |
| Lantana | Blood leaf | Purple coneflower |
| Lilac | Bud's | Rose moss |
| Locust | Candle larkspur | Saxia |
| Magnolia | Candytuft | Scarlet sage |
| Maple | Cinquefoil | Shrimp shell |
| Muck orange | Chrysanthemum | Stone crop |
| Mountain laurel | Colombine | Strawflower |
| Oak | Coleus | Sundrops |
| Pachistima | Cornopsis | Sunflower |
| Pine | Coral bells | Sweet pea |
| Pittosporum | Cosmos | Verbena |
| Podocarpus | Cypripedium | Vig. Spiderwort |
| Poplar | Dahlia | Violet |
| Privet | Delphinium | Wormwood |
| Redbud | Evening primrose | Zinnia |

DO NOT USE on the following plants:
Bottle Weed
Bottle Pink
Clematis
Crown
Geranium
Mossy geranium
Pansy
Phlox
Sweet William
T. anthera

CAUTION. Do not take internally. Avoid inhalation of dust. Avoid contact with eyes and skin. Wash after handling or use. Keep children and pets off treated areas until this material is washed into the soil and grass is dry. Do not graze treated areas or feed plant refuse to livestock. Avoid contamination of feed and foodstuffs. Do not store near seeds, fertilizer, or other pesticides.

NOTICE. Seller makes no warranty of any kind, express or implied, concerning effects of use of this product other than those specified on this label. Buyer, or User, accepts all responsibility for results due to misuse of this product.

USDA Reg. No. 2125-

Distributed in U.S.A. by
SCIENCE PRODUCTS COMPANY, Inc.
Chicago, Illinois

EDGING

For a weed-free border along walks, driveways, flower beds and ornamental gardens. Spray to wet thoroughly all plants in strips or areas desired to be free of all vegetation. Do not allow spray to contact desired plants.

LAWN RENOVATION

To kill off vegetation in a lawn area to be reseeded apply 1 gallon spray solution per 250 sq. ft. of area. Completely wet all vegetation to be killed. Bermuda grass and other deeprooted plants may require a second treatment. Reapply in 5 days for such areas. About 5 days after last treatment, rake away killed foliage. Water area well and seed at rate for new lawns. Rake seed into soil, mulch lightly with straw, and keep area well watered. Keep children and pets off treated areas until after first rain or sprinkling following treatment. Do not track from treated to untreated areas.

CAUTION. HARMFUL IF SWALLOWED. Avoid contact with skin. Avoid breathing spray mist. Wash hands after use. Avoid storage near feed or food products. Keep children and domestic animals off treated areas until this material has been washed into soil. Avoid contamination of Domestic or irrigation water. **ANTIDOTE.** If taken internally, induce vomiting and **CALL A PHYSICIAN AT ONCE.**

NOTICE. Seller makes no warranty of any kind, express or implied, concerning effects of use of this product other than those specified on this label. Buyer, or User, accepts all responsibility for results due to misuse of this product.

SCIENCE PRODUCTS COMPANY, Inc.
Chicago, Illinois

Science

Grass & Weed TOP-KILLER

**For LAWN EDGING and
LAWN RENOVATION
Also Kills Unwanted Vegetation**

**Kills TOP-GROWTH Only
Does Not Sterilize Soil**

| ACTIVE INGREDIENTS | |
|--|-------|
| Sodium Cacodylate | 12.6% |
| Dimethylarsinic Acid (Cacodylic Acid) | 2.2% |
| INERT INGREDIENTS | 85.2% |
| TOTAL 100.0% | |

Total Arsenic (as elemental) all in water soluble form ... 12.7%
Product contains 1.24 lbs. cacodylic acid equivalent per gallon

CAUTION: KEEP OUT OF REACH OF CHILDREN
See other cautions on back panels

**READ ENTIRE LABEL CAREFULLY BEFORE USE.
OBSERVE ALL CAUTIONS. USE ONLY AS DIRECTED.**

Grass & Weed TOP-KILLER is useful for general post-emergent weed control. It is a combination of herbicide and surfactant. It is non-selective in its action. Its phytotoxic properties are quickly inactivated on contact with the soil. Best results are obtained when applied to young, actively-growing weeds. It produces TOP-KILL only so repeat applications are needed for season-long weed control.

Grass & Weed TOP-KILLER is excellent for chemical edging of lawns, walks, driveways, flower beds, trees, shrubs, lawn ornaments, alongside buildings and fences. After use, clean equipment by flushing with water. Do not reuse empty container. Wash with water and detergent and discard in a safe place.

DIRECTIONS FOR USE

Thoroughly dissolve 5 1/3 fl. oz. (10 and 2/3 tablespoons) per gallon of water to make spray solution.

To TOP-KILL Vegetation

Along walks, driveways, fence lines, similar areas. Apply spray to wet all foliage thoroughly. On perennial weeds, repeat application as necessary.

CLEAN-UP around Trees, Shrubs, Evergreens

Direct spray onto vegetation to be controlled at base of trees and shrubs. Do not apply to or allow spray to contact foliage, green bark, graft unions, or broken bark when spraying around trees and shrubs.

USDA Reg No. 2125-56

Apr. 2, 1971
2125-56