

ACCEPTED
APR 2 1973
2125-58

VEGETABLES. Beans, cabbage, cantaloupes, carrots, cucumbers, onions, peppers, tomatoes. Controls aphids, bean beetles, cabbage worms, leaf miners, leafhoppers. Apply 3 gals. per 1,000 sq. ft. Cover leaves and fruits with a fine mist. Do not spray within 14 days of picking (1 day for tomatoes).

NUISANCE INSECTS. Outside areas only. Ants, boxelder bugs, chiggers, clover mites, earwigs, flies, roaches, spiders, ticks. Add 16 fl. oz. per 3 gallons water. Thoroughly spray window frames, doors, sills, screens, porch and patio walls and ceilings, foundations, walks, garbage cans, into cracks and other places where insects hide. Do not treat plants or animals with this dosage rate. Do not use indoors.

WARNING. May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes, clothing. Wash thoroughly after handling. Avoid contamination of feed, foodstuffs, food utensils. Do not use on household pets or humans. Keep out of reach of domestic animals. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. **COMBUSTIBLE MIXTURE** • Do not store or use near heat or open flame.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply when runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Break empty bottle; bury away from water supply, crops.

NOTICE. Buyer, or User, assumes all risks of use, storage or handling of this material not in strict accordance with directions given on this label.

Manufactured in U.S.A.

Distributed by

SCIENCE PRODUCTS COMPANY, Inc.
Chicago, Illinois

Covers 1,000 sq. ft.
of lawn area

Science

DIAZINON®

Spray

Versatile, Multi-Purpose Insect Spray
Controls Certain
LAWN, GARDEN, SOIL & NUISANCE INSECTS

Ants, Bermuda Mites, Chinch-bugs, Sod Webworms,
Earwigs, Roaches, Spiders, and certain others

ACTIVE INGREDIENTS

0,0-diethyl 0-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate	12.5 %
Aromatic Petroleum Derivative Solvent	72.6 %

INERT INGREDIENTS	14.9 %
-------------------	--------

TOTAL	100.0 %
-------	---------

DIAZINON, Reg. trademark of CIBA-GEIGY Corporation. U.S. Patent No. 2,754,243

WARNING: KEEP OUT OF REACH OF CHILDREN
See other cautions on back panels

FUNGICIDE FOR ECONOMIC POISON REGISTERED UNDER FEDERAL TO ATTACHED COMMENTS

READ ENTIRE LABEL CAREFULLY BEFORE USE.
OBSERVE ALL CAUTIONS. USE ONLY AS DIRECTED.

GENERAL. Unless otherwise directed mix 4 teaspoons this product in 1 gal. water and apply full amount evenly over a 40 sq. ft. area. For lawns, vegetables measure size of area to be sprayed (20' x 50' = 1,000 sq. ft.). Then mix proper amount of solution and spray as directed. Use coarse spray for lawns, fine mist spray for plants, trees, shrubs. Application remains effective for an extended period of time. Flush sprayer with clean water after use. Repeat as needed.

LAWNS. Merion, Bluegrass, Fescue, Dichondra, St. Augustine, Bent, Bermuda, Lippia varieties. Controls chinch-bugs, sod webworms (lawn moths), Bermuda mites, ants, ticks, earwigs, billbugs, chiggers, clover mites, leafhoppers, cutworms, armyworms, digger wasps. Thoroughly wet grass a few hours before applying; for billbugs water lawn after treatment. For millipedes use 8 teaspoons per gallon for each 40 sq. ft. of lawn. Do not permit children or pets to go onto the sprayed grass until the spray has completely dried.

ORNAMENTALS. Annual and perennial flowers, roses, evergreens, trees, shrubs. Controls aphids, bagworms, leafhoppers, leaf miners, mites, red spiders, scale crawlers, tent caterpillars, thrips, webworms, whiteflies. Spray top and underside of leaves and branches to dripping point. On trees, shrubs try to penetrate dense foliage. Do not use on ferns, poinsettias, hibiscus, gardenias, orchids.

FRUITS. Apples, oranges, peaches, plums, grapes, strawberries. Controls aphids, mites, codling moth, leaf rollers, grape leaf folder, scale crawlers. Start spraying after blooming and repeat at 10 to 14 day intervals if needed. Apply as full cover spray. Do not spray fruit within 21 days of picking.

EPA Reg. No. 2125-58-AA

VEGETABLES. Bear 3, cabbage, cantaloupes, carrots, cucumbers, onions, peppers, tomatoes. Controls aphids, bean beetles, cabbage worms, leaf miners, leafhoppers. Apply 3 gals. per 1,000 sq. ft. Cover leaves and fruits with a fine mist. Do not spray within 14 days of picking (1 day for tomatoes).

NUISANCE PESTS. Outside areas only. Ants, boxelder bugs, chiggers, clover mites, earwigs, flies, roaches, spiders, ticks. Add 16 fl. oz. per 3 gallons water. Thoroughly spray window frames, doors, sills, screens, porch and patio walls and ceilings, foundations, walks, garbage cans, into cracks and other places where insects hide. Do not treat plants or animals with this dosage rate. **Do not use indoors.**

WARNING. May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes, clothing. Wash thoroughly after handling. Avoid contamination of feed, foodstuffs, food utensils. Do not use on household pets or humans. Keep out of reach of domestic animals. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. **COMBUSTIBLE MIXTURE** • Do not store or use near heat or open flame.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply when runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Break empty bottle; bury away from water supply, crops.

NOTICE. Buyer, or User, assumes all risks of use, storage or handling of this material not in strict accordance with directions given on this label.

Manufactured in U.S.A.

Distributed by

SCIENCE PRODUCTS COMPANY, Inc.
Chicago, Illinois

Covers 1,000 sq. ft.
of lawn area



Net Contents
1 Pint

DIAZINON®

Spray

Versatile, Multi-Purpose Insect Spray
Controls Certain
LAWN, GARDEN, SOIL & NUISANCE INSECTS

Ants, Bermuda Mites, Chinch-bugs, Sod Webworms,
Earwigs, Roaches, Spiders, and certain others

ACTIVE INGREDIENTS

0,0-diethyl 0-(2-isopropyl-4-methyl-
6-pyrimidinyl) phosphorothioate 12.5 %

Aromatic Petroleum Derivative Solvent 72.6 %

INERT INGREDIENTS 14.9 %

TOTAL 100.0 %

®DIAZINON, Reg. trademark of CIBA-GEIGY Corporation. U.S. Patent No. 2,754,243

WARNING: KEEP OUT OF REACH OF CHILDREN
See other cautions on back panels

READ ENTIRE LABEL CAREFULLY BEFORE USE.
OBSERVE ALL CAUTIONS. USE ONLY AS DIRECTED.

GENERAL. Unless otherwise directed mix 4 teaspoons this product in 1 gal. water and apply full amount evenly over a 40 sq. ft. area. For lawns, vegetables measure size of area to be sprayed (20' x 50' = 1,000 sq. ft.). Then mix proper amount of solution and spray as directed. Use coarse spray for lawns, fine mist spray for plants, trees, shrubs. Application remains effective for an extended period of time. Flush sprayer with clean water after use. Repeat as needed.

LAWNS. Merion, Bluegrass, Fescue, Dichondra, St. Augustine, Bent, Bermuda, Lippia varieties. Controls chinch-bugs, sod webworms (lawn moths), Bermuda mites, ants, ticks, earwigs, billbugs, chiggers, clover mites, leafhoppers, cutworms, armyworms, digger wasps. Thoroughly wet grass a few hours before applying; for billbugs water lawn after treatment. For millipedes use 8 teaspoons per gallon for each 40 sq. ft. of lawn. Do not permit children or pets to go onto the sprayed grass until the spray has completely dried.

ORNAMENTALS. Annual and perennial flowers, roses, evergreens, trees, shrubs. Controls aphids, bagworms, leafhoppers, leaf miners, mites, red spiders, scale crawlers, tent caterpillars, thrips, webworms, whiteflies. Spray top and underside of leaves and branches to dripping point. On trees, shrubs try to penetrate dense foliage. Do not use on ferns, poinsettias, hibiscus, gardenias, orchids.

FRUITS. Apples, oranges, peaches, plums, grapes, strawberries. Controls aphids, mites, codling moth, leaf rollers, grape leaf folder, scale crawlers. Start spraying after blooming and repeat at 10 to 14 day intervals if needed. Apply as full cover spray. Do not spray fruit within 21 days of picking.

EPA Reg. No. 2125-58-AA

Mar. 7, 1973

2125-58