

Science

ZINEB

Garden Fungicide

Controls certain diseases of
FLOWERS, VEGETABLES, FRUITS,
LAWNS and ORNAMENTALS

Anthracnose, Early Blight, Late
Blight, Downy Mildew, Rust and
certain other diseases specified
under Directions For Use.

CAUTION: KEEP OUT OF REACH OF CHILDREN

See other cautions on back panel.

READ ENTIRE LABEL CAREFULLY BEFORE USE. OBSERVE ALL CAUTIONS. USE ONLY AS DIRECTED.

GENERAL DIRECTIONS FOR USE

Thorough and uniform coverage is absolutely essential for disease control. Apply spray mix to both under and upper surfaces of foliage as well as entire plant.

VEGETABLE CROPS

Use one tablespoonful per gallon of water unless otherwise shown. Start applications at first sign of, or prior to appearance of, disease and repeat as directed. Do not apply within 7 days of harvest on leafy vegetables, or on root crops if tops are to be used as food or feed.

BEANS. Anthracnose, Downy mildew, Rust. Begin when first blossoms appear and repeat at 7 to 14 day intervals. Do not apply within 7 days of harvest.

CARROTS. Alternaria leaf blight, Cercospora leaf spot.

KALE, COLLARDS MUSTARD GREENS, SWISS CHARD. Cercospora leaf spot, Downy mildew.

CANTALOUPE, CUCUMBERS, MELONS, PUMPKINS, SQUASH. Alternaria blight, Anthracnose, Downy mildew, Gummy stem blight. Apply at 7 to 14 day intervals. Cover underside of leaves thoroughly.

LETTUCE. Downy mildew. Apply at weekly intervals to within 5 days of harvest for head lettuce; 7 days for leaf lettuce.

TOMATOES. Anthracnose, Cladosporium gray mold, early and late blight, Stemphylium and Septoria leaf spots. Apply at 7 to 10 day intervals; oftener if disease is severe.

SPINACH. Downy mildew (Blue mold), White rust.

FLOWERS AND ORNAMENTALS

Use 1 level tablespoonful per gallon of water. Spray plants thoroughly.

AZALEAS, CAMELLIAS. Petal blight. Spray 2 to 3 times per week while flowers are opening. Direct spray into flowers. Spray ground under bushes thoroughly.

GLADIOLUS. Botrytis blight, Curvularia, Stemphylium. Spray at weekly intervals, or oftener under severe disease conditions, from early stages until spikes begin to open.

SNAPDRAGONS, CHRYSANTHEMUMS, CARNATIONS, HOLLYHOCKS. Anthracnose, Leaf spot, Rust. Apply at weekly to 10 day intervals as soon as disease conditions threaten.

ROSES. Black spot. Apply when foliage first appears. Repeat at weekly intervals.

FRUITS

Mix amounts of ZINEB indicated in 1 gallon of water to make spray solution. In home plantings apply at 1 to 2

week intervals beginning before bloom. Spray plants thoroughly.

APPLES. 1 tablespoon. Apple blotch, Bitter rot, Black rot, Botryosphaeria fruit rot, Brooks spot, Cedar rust, Fly speck, Frog-eye leaf spot, Quince rust, Scab, Sooty blotch. Begin applications in delayed dormant and continued through cover sprays.

APPLES, PEARS. 7/8 tablespoons. Fire blight. Apply at 2 to 3 day intervals when new blooms are opening and weather is wet. Use sanitary pruning practices. Do not apply to pears within 7 days of harvest.

PEACHES. 3 teaspoons. **APRICOTS.** 4 teaspoons. Coryneum blight or shot hole, Peach leaf curl. Apply in post-harvest or dormant period. Do not apply after petal fall on apricots or within 30 days of harvest on peaches.

STRAWBERRIES. 4 teaspoons. Leaf spot. Begin when new growth starts in Spring and repeat as necessary. Leaf scorch: apply in new beds in late summer using 2 to 3 weekly applications. Do not apply within 7 days of harvest.

SOUR CHERRIES. 2 teaspoons. Leaf spot or shot hole. Apply at petal fall, and again immediately after harvest. Do not apply within 7 days of harvest.

GRAPES. 3 teaspoons. Black rot, Downy mildew, Ripe rot. Apply just before bloom, just after bloom, 10 days after bloom, and 2 to 3 weeks later. Do not apply within 7 days of harvest.

LAWNS AND TURFS

Use 2 oz. (8 slightly rounded tablespoons) ZINEB in 3 to 5 gallons water and apply full amount uniformly to 1,000 sq. ft. of lawn area. Or use 2 1/2 teaspoons ZINEB in 1 gallon water for 100 sq. ft. area. Begin applications as soon as diseases appear.

For Fading-out (Curvularia), Helminthosporium diseases, Rust, and Slime molds repeat every 7 to 14 days until disease threat is past. For grease spot or Pythium blight repeat every 5 to 7 days when disease is mild or every 3 to 4 days when disease is severe. Continue applications until disease threat is past. Consult State Agricultural Experiment Station or State Extension Specialist for proper timing and application in your area.

CAUTION. May cause irritation of nose, throat, and skin. Do not breathe dust or spray mist. In case of skin contact, wash with water; for eyes, wash thoroughly with water and get medical attention. Keep container closed. Store in a dry place. Keep away from fire and sparks. Do not reuse this container. Destroy it when empty.

NOTICE. Seller makes no warranty of any kind, express or implied, concerning effects of use of this product other than those specified on this label. Buyer, or User, accepts all responsibility for results due to misuse of this product.

SCIENCE PRODUCTS COMPANY, INC.

Nov. 10, 1971

