

LAWNS. Use 1/2 pint (8 fl. oz.) in 15-20 gals. water and apply evenly to 500 sq. ft. of lawn area. Use garden hose sprayer or pressure-type equipment and apply full water volume to insure good penetration of turf. For ants, chinch-bugs, earwigs, fall armyworm, fleas, leafhoppers, millipedes, mosquitoes, sod webworm (lawn moth). For best results mow lawn and apply immediately after rain or watering. Repeat 2 to 3 weeks later, if necessary. In Florida use 10 oz. for chinch-bug control.

CAUTION. HARMFUL IF SWALLOWED OR INHALED. Avoid prolonged breathing of spray. Do not take internally. Avoid prolonged or repeated contact with skin. Wear long-sleeved clothing and clean after spraying. Wash thoroughly after handling. Wash hands and face before eating. To avoid injury to tender foliage do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not reuse empty container. Destroy by breaking. Discard in a safe place. Do not use on Boston Ivy.

Avoid contamination of food, feed, water supplies, streams and ponds.

NOTE FOR PHYSICIAN. Carbaryl is a moderately reversible, cholinesterase inhibitor. Atropine is antidotal.

NOTICE. Buyer, or User, assumes all risks of use, storage, or handling of this material not in strict accordance with the directions given on this label.

Manufactured in U.S.A. for

SCIENCE PRODUCTS COMPANY, Inc.
Chicago, Illinois 60646

Science SEVIN[®] Liquid

Kills
CHINCH-BUGS, JAPANESE BEETLES, BAGWORMS,
EARWIGS, LAWN MOTHS (sod Webworm),
and many other insects on

Flowers — Vegetables — Fruit
Lawns — Shrubs — Ornamentals

ACTIVE INGREDIENT
Carbaryl (1-naphthyl N-methyl
carbamate)
INERT INGREDIENTS
TOTAL 100.0%
Contains 2 lbs. SEVIN[®] per gallon
SEVIN, registered trade mark of Union
Carbide Corporation for carbaryl insecticide
U.S. Pat. Nos. 2,903,478; 3,009,855

Net Contents: 1 Pint

CAUTION: KEEP OUT OF REACH OF CHILDREN
See other cautions on back panels

READ ENTIRE LABEL CAREFULLY BEFORE USE.
OBSERVE ALL CAUTIONS. USE ONLY AS DIRECTED.

DIRECTIONS FOR USE (Shake well before using.)

Unless otherwise directed mix 4 teaspoons this product per gallon of water. Spray entire plant thoroughly when insects or their damage first appear. Repeat at 7 to 10 day intervals, as needed. Keep spray mixture well-agitated.

ORNAMENTALS. Flowers, shrubs, trees. For apple aphid, bagworm, birch leaf miner, blister beetle, boxelder bug, boxwood leaf miner, elm leaf aphid, elm leaf beetle, flea beetle, gypsy moth, Japanese beetle, lace bug, leafhopper, leaf roller, mealy bug, mimosa webworm, oak leaf miner, orange tortrix, periodical cicada (17 year locust), plant bug, puss caterpillar, psyllids, rose aphid, rose slug, scale insects, tent caterpillar, thorn bug, exposed thrip, willow leaf beetle.

VEGETABLES. Use 4 tsps. per gal. water on young plants. 8 teaspoons on mature plants for corn earworm, European corn borer, fall armyworm, flea beetles, leafhoppers, cucumber beetles, pickleworm, stink bugs, tomato fruitworm, tomato hornworm on corn, beans, (green, lima, snap), eggplant, peppers, peas; on cucumbers, melons, pumpkins, squash use only 4 tsps. per gal. Do not use on watermelons in Florida. Application to tassel region of corn during pollen shed period will seriously reduce bee population. Do not apply to listed crops within 1 day of harvest.

FRUIT and CITRUS TREES. For codling moth, cucumber beetle, fruit tree leaf roller, lesser peach tree borer, plum curculio, red banded leaf roller, apple maggot, bagworm, citrus cutworm, orange tortrix, Forbes scale, San Jose scale, on apples, pears, citrus, apricot, nectarine, peaches, plums, cherries, and shade trees. Do not apply within 5 days of harvest on citrus trees, apricot, nectarine, nor within 1 day of harvest on other fruit trees listed.

E.P.A. Est. 2125-II-1
EPA Reg. No. 2125-61-AA

2125-61

05/06/1976

1/1