

when rain or excessive humidity is expected during the next 2 days. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not reuse empty container. Destroy by breaking. Discard in a safe place.

Avoid contamination of food, feed, water supplies, streams and ponds

NOTE FOR PHYSICIAN: Carabaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

NOTICE: Buyer, or User, assumes all risks of use, storage or handling of this material not in strict accordance with the directions given on this label.

Manufactured in U.S.A. for

SCIENCE PRODUCTS COMPANY, Inc.
Chicago, Illinois

**EARWIGS, LAWN MOTHS (sod Webworms),
and many other insects on**

**Flowers — Vegetables — Fruits
Lawns — Shrubs — Ornamental**

ACTIVE INGREDIENT

Carbaryl (1-naphthyl N-methyl- carbamate)	24.4%
INERT INGREDIENTS	75.6%
TOTAL	100.0%

Contains 2 lbs. SEVIN* per gallon
*SEVIN, registered trade mark of Union Carbide Corporation for carbaryl insecticide
U.S. Pat. Nos. 2,903,478 3,009,855

Net Contents: 1 Pint

CAUTION: KEEP OUT OF REACH OF CHILDREN
See other cautions on back panels

AC
9
12

50W

Kill
JAPANESE BEETLES
GRASSHOPPER
17 YEAR LOCUST

many other garden insects
on
FLOWERS — FRUITS — VEGETABLES
LAWNS—SHRUBS—ORNAMENTALS

ACTIVE INGREDIENTS
Carbaryl—(1-Naphthyl Methylcarbamate) 50%
INERT INGREDIENTS 50%

U.S. Patent Nos. 2,903,478; 3,009,855.

*SEVIN is the registered trademark of Union Carbide Corporation for carbaryl insecticide.

LAWNS

For lawns, use 1/2 lb. per 1000 sq. ft. of lawn area. Apply in early morning or late afternoon. Do not water for 24 hours after application.

VEGETABLES

For most vegetables, use 1/2 lb. per 1000 sq. ft. of area. For beans, use 1 lb. per 1000 sq. ft. of area. For corn, use 1/2 lb. per 1000 sq. ft. of area.

BEANS. For Mexican bean beetle, use 1 lb. per 1000 sq. ft. of area. For other insects, use 1/2 lb. per 1000 sq. ft. of area.

CORN. Use 1/2 lb. per 1000 sq. ft. of area. For European cornborer, use 1 lb. per 1000 sq. ft. of area.

CUCUMBERS, SQUASH, PUMPKINS, MELONS. Use 1/2 lb. per 1000 sq. ft. of area. For cucumber beetle, use 1 lb. per 1000 sq. ft. of area.

TOMATOES, POTATOES, EGGPLANT, PEPPERS. Use 1/2 lb. per 1000 sq. ft. of area. For Colorado potato beetle, use 1 lb. per 1000 sq. ft. of area.

OKRA. Use 1/2 lb. per 1000 sq. ft. of area. For okra beetle, use 1 lb. per 1000 sq. ft. of area.

CABBAGE, BROCCOLI, CAULIFLOWER, BRUSSELS SPROUTS, KOHLRABI. Use 1/2 lb. per 1000 sq. ft. of area. For cabbage looper, use 1 lb. per 1000 sq. ft. of area.

FRUITS. Use 1/2 lb. per 1000 sq. ft. of area. For apple maggot, use 1 lb. per 1000 sq. ft. of area.

APRICOTS, PEACHES, NECTARINES. Use 1/2 lb. per 1000 sq. ft. of area. For peach twig borer, use 1 lb. per 1000 sq. ft. of area.

CITRUS. Use 1/2 lb. per 1000 sq. ft. of area. For citrus red scale, use 1 lb. per 1000 sq. ft. of area.

GRAPES. Use 1/2 lb. per 1000 sq. ft. of area. For grape leafhopper, use 1 lb. per 1000 sq. ft. of area.

BLUEBERRIES. Use 1/2 lb. per 1000 sq. ft. of area. For blueberry leaf miner, use 1 lb. per 1000 sq. ft. of area.

CRANBERRIES, STRAWBERRIES. Use 1/2 lb. per 1000 sq. ft. of area. For strawberry root weevil, use 1 lb. per 1000 sq. ft. of area.

CHERRIES, PLUMS, PRUNES. Use 1/2 lb. per 1000 sq. ft. of area. For cherry bark beetle, use 1 lb. per 1000 sq. ft. of area.

CAUTION: HARMFUL IF SWALLOWED OR INHALED.

NOTICE

SCIENCE PRODUCTS COMPANY, Inc.
Chicago, Illinois
EPA Reg. No. 2125-63 AA

50W

LEAF LOCUS

**many other garden insects
on
FLOWERS — FRUITS — VEGETABLES
LAWNS—SHRUBS—ORNAMENTALS**

ACTIVE INGREDIENTS
Carbaryl—(1-Naphthyl Methylcarbamate) .50%
INERT INGREDIENTS50%

U.S. Patent Nos. 2,903,478; 3,009,855.

CAUTION: KEEP OUT OF REACH OF CHILDREN
See other cautions on back panels