

ALL OTHER LISTED INSECTS IN  
1. WAF - FROGS - TOADS -  
2. WAF - GRAPES - CUCUMBERS

ACTIVE INGREDIENTS:  
Dibutyl (2-ethylhexyl) sebacate, ----- 90.05  
INERT INGREDIENTS: ----- 9.95  
100.00  
EPA Reg. No. 111-6; EPA Act. No. 2127-11-1

SAVING is the registered trademark of the Davison Corporation for  
dibutyl sebacate.  
U.S. Pat. Nos. 2,813,415; 3,007,811.

ALL INFORMATION SUBJECT TO CHANGE

CAUTION

See Side Label for Additional Precautionary Statements.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by  
touching back of throat with finger. Call physician or Poison Control  
Center. Do not induce vomiting or give anything by mouth to an unconscious  
person. IF INHALED: Remove victim to fresh air. IF IN EYES: Flush  
with plenty of water. IF ON SKIN: Remove contaminated clothing. Wash  
affected area with soap and water. Shower completely before eating.  
NOTE: DO NOT USE THIS PRODUCT IN A MANNER, particularly in the absence of  
inhibition, intended or in violation of applicable regulations.

DAVISON CHEMICAL & FERTILIZER DIVISION  
P.O. BOX 1000  
LANCASTER, PENNSYLVANIA 17604

DAVISON CHEMICAL & FERTILIZER DIVISION

PRECAUTIONS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS--CAUTION: Harmful if inhaled or swallowed. Do not breath dust or spray mist. Avoid contact with skin, eyes, or clothing. Do not take internally. Wear long-sleeves. Change clothing daily. Wash hands and face before eating. Wash thoroughly after handling. Keep children and pets off treated area until spray has dried. ENVIRONMENTAL HAZARDS: Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from treated area. Do not contaminate water by cleaning of equipment or disposal of waste. BEE CAUTION: MAY KILL BEEKEEPERS IN SUBSTANTIAL NUMBERS. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not use when value of bees as pollinators is more important than insect control. Warn beekeepers to locate hives beyond bee flight range until one week after application, or take equally effective precautions.

DEFINITION OF TERM

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Store in closed, original container in a locked area away from children and domestic animals. For disposal, securely wrap closed container in several layers of newspaper and place in trash collection. Do not reuse container.

Use 2 level tablespoons in 1 gallon water, unless otherwise directed. Mix proper amount in a little water to make a paste, then add balance of water to make a gallon of spray mixture. Make first application when insects or their damage first appear. Repeat applications at 7 to 10 day intervals to maintain effective control, unless otherwise directed. Keep spray mixture well-agitated.

**ORNAMENTALS**

**ROSES, OTHER FLOWERS, SHRUBS, SHADE TREES.** To control Japanese beetles, apple aphid, bagworm, birch leaf miner, elm leaf aphid and beetle, flea beetles, grape moth, June beetles, lace bugs, leaf hoppers, leaf rollers, mealy bugs, mimosa webworms, oak leaf miners, orange tortrix, periodical cicada (17 year locust), plant bugs, puss caterpillars, psyllids, rose aphid and slug, scale insects, tent caterpillars, thorn bug, exposed thrips, willow leaf beetle. **DO NOT USE ON BOSTON IVY.**

**LAWNS**

To control chinch bugs, ants, earwigs, fall armyworms, fleas, leaf hoppers, millipedes, mosquitoes, sod webworms (lawn moths). Use 1 level cupful in 15 to 20 gallons water and apply evenly to 500 sq. ft. of established lawn area.

**VEGETABLES**

Can be used on day of harvest unless otherwise indicated. **BEANS.** For Mexican bean beetle, bean leaf beetle, Western bean cutworm, cucumber beetles, flea beetles, Japanese beetles, leafhoppers, velvet bean caterpillar, armyworms, corn earworm, stinkbugs, tarnished plant bug. In California use 4 tablespoons per gallon for corn earworm, lima bean pod borer, Lygus and stink bugs.

**CARROTS, LETTUCE.** 2 to 4 tablespoons. Do not use on head lettuce within 3 days of harvest nor on leaf lettuce within 14 days of harvest. For flea beetles, leafhoppers, harlequin bug, six-spotted leafhopper (aster yellows vector), armyworms, cabbage worm, corn earworm, aphids, bugs, tarnished plant bug, stink bugs. For corn earworm on lettuce, use every 3 to 7 days after heads form.

**CORN.** Use 2 to 4 tablespoons. For corn rootworm adults, fall armyworm, flea beetles, Japanese beetles, sap beetles, leafhoppers. Corn earworm, apply to silks as soon as they appear. Make 3 to 4 applications at 2-3 day intervals. Reapply immediately after rain. Application to the tassels region of corn during the pollen shed period will seriously reduce bee population. European corn borer, apply thoroughly into whorls when plant is knee-high.

**CUCUMBERS, SQUASH, PUMPKINS, MELONS.** For pickleworm, melon worm, cucumber beetles, flea beetles, leafhoppers, squash bugs. Do not use on watermelons in Florida.

**TOMATOES, POTATOES, EGGPLANT, PEPPERS.** Use 2 to 4 tablespoons. For Colorado potato beetle, flea beetles, leafhoppers, European corn borer, fall armyworm, lace bug, tomato fruitworm, tomato hornworm, tarnished plant bug, stink bugs. **OKRA.** For corn earworm and stinkbugs.

**CABBAGE, BROCCOLI, CAULIFLOWER, BRUSSELS SPROUTS, KOHLRABI.** Use 2 to 4 tablespoons. Do not use within 3 days of harvest. For flea beetles, harlequin bug, armyworms, cabbage worm, corn earworm.

**FRUITS**

Use up to 1 day before harvest unless otherwise directed. Apply at petal fall and every 10-14 days to maintain control.

**APPLES, PEARS.** West of Rockies: For codling moth, apple sucker, green and woolly apple aroid, bagworm, California pear slug, apple and pear rust mites, pear leaf blister mite, eye-spotted bud moth, green fruitworm, Lygus bug, orange tortrix, pear psylla, tentiform leaf mite, Lecanium oyster shell and San Jose scale. Delay use until 30 days after full bloom to avoid possibility of fruit thinning. East of Rockies: 1 tablespoon for codling moth, apple mealybug, green apple aphid, white apple leafhopper. 2 tablespoons for apple maggot, bagworm, Eastern tent caterpillar, European apple saw fly, eye-spotted bud moth, red-banded and fruit tree leaf rollers, green fruitworm, Japanese beetle, pear psylla, periodical cicada (17 year locust), plum curculio, sawy and woolly apple aphids, apple and pear rust mites, pear leaf blister mites, tarnished plant bug, tentiform leaf miner, Forbes, Lecanium, oyster shell and San Jose scales.

**APRICOTS, PEACHES, NECTARINES.** For Oriental fruit moth, codling moth, cucumber beetles, European earwig, Japanese beetle, Pandemis moth, June beetles, lesser peach tree borer, orange tortrix, peach twig borer, periodical cicada, plum curculio, red-banded and fruit-tree leaf rollers, Plectyotoma flavendana, tarnished plant bug, tussock moth, Lecanium, San Jose and olive scales. Do not use within 3 days of harvest on apricots, nectarines.

**CITRUS.** For citrus cutworm, California orange dog, fruit tree leaf roller, orange tortrix, Western tussock moth, black scale, brown soft scale, California red scale, citrus scale, yellow scale. Do not use within 5 days of harvest.

**GRAPES.** Can be used on day of harvest. For European fruit Lecanium grape leaf folder, grape leafhopper, grape leaf skeletonizer, grape berry moth, Japanese beetle, June beetle, red-banded leaf roller.

**BLUEBERRIES.** Can be used on day of harvest. For blueberry maggot, cherry and cranberry fruitworms, European fruit Lecanium, Japanese beetle.

**CRANBERRIES, STRAWBERRIES.** Use up to 1 day before harvest. For cutworms, fireworms, fruitworms, Japanese beetle, leafhoppers, meadow spittlebug, strawberry leaf-roller, strawberry weevil.

**CHERRIES, PLUMS, PRUNES.** Use up to 1 day before harvest. For black cherry aphid, mealy plum aphid, cherry fruit fly, cherry fruitworm, eye-spotted bud moth, red-banded and fruit-tree leafroller, Japanese beetle, lesser peach twig borer, plum curculio, prune leafhopper, brown soft scale, Forbes, Lecanium, oyster shell and San Jose scales, peach twig borer, Eastern tent caterpillar, codling moth, orange tortrix, tussock moth.

**CAUTION: HARMFUL IF SWALLOWED OR INHALED.** Do not take internally. Avoid breathing dust or spray mist. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of feed or foodstuffs. Keep children and pets off treated areas until this material is washed into the soil and grass is dry. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from the Cooperative Agricultural Extension Service. Destroy empty container and bury it in a safe place.

**LIMITED WARRANTY: BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.**