

**DIRECTIONS FOR USE**

Begin application when disease first appears. Repeat at 10-14 day intervals throughout growing season, or as long as needed.

**ROSES** (Field & Greenhouse). For Black Spot, Powdery Mildew. Mix 1 tablespoon in 2 gallons water. Shorten spray interval during humid, rainy weather.


**FLOWERS, ORNAMENTALS** (Field & Greenhouse). For Powdery Mildew, Betrytis Grey Mold. Mix 1 tablespoon in 2 gallons water.

**LAWNS.** Dollar Spot (*Sclerotinia homeocarpa*), Large Brown Patch (*Rhizoctonia solani*). Apply at rate of 2 oz. (12 tablespoons) per 1,000 sq. ft. in enough water to obtain thorough coverage, usually 2 to 5 gals. per 1,000 sq. ft. Apply every 5 to 7 days when conditions especially favor disease. Do not graze or feed clippings from treated areas to livestock.

EPA Reg. No. 2125-64-AA

Distributed by

**SCIENCE PRODUCTS COMPANY, Inc.**  
Chicago, Illinois



# BENOMYL

BLACK SPOT &  
POWDERY MILDEW

on ROSES  
Flowers, Ornamentals  
and for certain  
LAWN DISEASES

Net Wt. 2 ozs.

**CAUTION:** KEEP OUT OF REACH OF CHILDREN  
MAY IRRITATE EYES, NOSE, THROAT, SKIN  
See other cautions on side panel

**CAUTION.** Avoid breathing dust. Avoid contact with skin, eyes, clothing. Wash thoroughly after use. Keep away from fire or sparks. In case of contact flush skin or eyes with plenty of water; for eyes, get medical attention.

**IMPORTANT.** Never allow BENOMYL to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of BENOMYL as a fungicide. Keep container closed when not in use. Do not re-use container. Bury when empty.

BENOMYL is highly effective for control of certain fungus diseases of roses, flowers, ornamentals, and lawns providing both curative (eradicator) action and protective action. It is a wettable powder to be mixed with water for application as a spray. Addition of a surfactant to the spray mix enhances curative action of BENOMYL; improves distribution of the spray on hard-to-wet plants such as roses (field & greenhouse).

**ACTIVE INGREDIENT**

Benomyl [Methyl (1-butylcarbamoyl)  
-2-benzimidazolecarbamate] .....50%  
INERT INGREDIENTS .....50%

U.S. Pat. Pending

3-21-72  
2125-64