



SCIENCE 5% DIAZINON® DUST

Kills ANTS and Other Lawn and Soil Insects. Also Foliage Protection for Flowers, Ornamentals and Certain Fruits and Vegetables.

ACTIVE INGREDIENT:

Diethyl O-(2-isopropyl-6-methyl-4-pyridinyl) phosphorothioate

Diethyl O-(2-isopropyl-6-methyl-4-pyridinyl) phosphorothioate	-----5.0%
INERT INGREDIENTS:	-----95.0%
TOTAL	-----100.0%

EPA Reg. No. 2125-65

EPA Est. No. 2125-IL-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT


IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Call physician or Poison Control Center. Do not induce vomiting or give anything by mouth to an unconscious person. IF IN EYES: Flush with plenty of water. Get medical attention. IF INHALED: Remove victim to fresh air. IF ON SKIN: Remove contaminated clothing. Wash affected area with soap and water. Launder clothing before reusing.

See Side Panel For Additional Precautionary Statements.

SCIENCE PRODUCTS DIVISION
MILLER CHEMICAL & FERTILIZER CORPORATION
Hanover, Pennsylvania 17331

NET WEIGHT: 1 POUND
4 POUNDS
8 POUNDS

ACCEPTED
DEC 23 1983
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 2125-65



BEST DOCUMENT AVAILABLE

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS--CAUTION: Harmful if swallowed. May be absorbed through the skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Do not contaminate water by cleaning of equipment or disposal of waste. Wash thoroughly after handling. Do not use on humans or household pets. Water lawns after application. Keep children and pets off treated area until dry. ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and other wildlife. Birds feeding on treated area may be killed. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of waste. Do not apply when weather conditions favor drift from treatment area. This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Store this product in its closed, original container in a locked area away from children and domestic animals. Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. For disposal, securely wrap original container in several layers of newspaper and discard in trash, or at an approved waste disposal facility.

BEST DOCUMENT AVAILABLE

GENERAL DIRECTIONS: When used on their own, they are most effective when applied to plants in the early stages of their growth. For best results, use the following directions:

VEGETABLES (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

FRUITS (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

ORNAMENTALS (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

SPINACH (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

POTATOES (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

APPLES (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

PLUMS (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

PEACHES (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

STRAWBERRIES (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

SOFT FLOWERS (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

CUTVEGETABLES (10):

Beans	Corn	Corn
Beets	Cauliflower	Cucumbers
Cabbage	Carrots	Radishes
Carrots	Celery	Spinach

GENERAL DIRECTIONS: When used on their own, they are most effective when applied to plants in the early stages of their growth. For best results, use the following directions:

VEGETABLES (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

FRUITS (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

ORNAMENTALS (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

SPINACH (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

POTATOES (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

APPLES (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

PLUMS (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

PEACHES (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

STRAWBERRIES (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

SOFT FLOWERS (10): Apply 1/2 cup of the concentrate to 100 sq. ft. of plants in the early stages of their growth. Repeat application as needed to control insects.

BEST DOCUMENT AVAILABLE

