ACCEPTED

This container is a

Refiliable SQUEEZE-DUSTER.

Lift hinged cap. Squeeze container to apply dust Refill through bottom closure.

READ ENTIRE LABEL THOROUGHLY BEFORE USE. USE ONLY AS DIRECTED, UNDICHEL ALL REPLANTED ACT OBSERVE CAUTION AND LIMITATION STATEMENTS CAREFULLY. FOR ECONOMIC POISON ADDITION

DIRECTIONS FOR USE

When insects or diseases first appear or threaten apply this dust uniformly-teentire plant including undersides of foliage. Repeat at 7-10 day intervals; more frequently during periods of prolonged rainfall or high humidity. Do not apply within number of days of harvest shown in parentheses after each plant listed. Do not apply in excess.

TOMATOES (5). Colorado Potato Beetle, Climbing Cutworm, Flea Beetle, Leafhopper, European Corn Borer, Fall Armyworm, Lace Bug, Pinworm, Stink Bug, Tomato Fruitworm, Tomato Hornworm, Tarnished Plant Bug; also Anthrecnose, Early and Late Blight, Gray Mold (Cladosporlum), Gray Leaf Spot (Stemphyllum), Nail Head Spot, Septoria Leaf Spot, Southern Blight (Scierotium).

BEANS (7). Green, Lima, Snap. Armyworm, Bean Leaf Beetle, Bean Leaf Roller, Corn Earworm, Cucumber Beetle, Cutworm, Flea Beetle, Japanese Beetle, Leafhopper, Lima Bean Pod Borer, Lygus Bug, Mexican Bean beetle, Stink Bug, Tarnished Plant Bug, Velvetbean Caterpillar, Western Bean Cutworm; also Anthracnose, Downy Mildew, Rust.

BROCCOLI (7), BRUSSELS SPROUTS (7), CABBAGE (7), CAULIFLOWER (7), CHINESE CABBAGE (14), COLLARDS (14), KÁLE (14), KOHLRABI (3), MUSTARD GREENS (14), RADISHES (3), TURNIPS: Roots (3), Tops (14). Armyworm, Corn Earworm, Diamondback Moth, Flex Beetle, Harlequin Bug, Imported Cabbageworm, Stink Bug, Tarnished Plant Bug; also Downy Mildew, Leaf Spots. Do not apply to kohlrabi after edible parts are half-grown.

CARROTS (7). Fies Beetle, Lesthopper, Six-Spotted Lesthopper, Spittlebug, Sunk bug; also Early Blight, Late Blight.

CORN (0). Corn Earworm (apply to silks as soon as they appear; make 3 to 4 applications at 2-3 day intervals as needed; re-apply immediately after rain), Corn Rootworm Adult, European Corn Borer (apply thoroughly into whorls when plants are knee-high; repeat weekly until tassels emerge), Fall Armyworm, Flea Beetle, Japanese Beetle, Leafhopper, Sap Beetle; Leaf Blight, Puccinia Rust. Application to tassel region of corn during pollen shed period will seriously reduce bee population. Do not feed treated forage or husks to dairy animals or animals being finished for slaughter.

CLCUMBERS (5), MELONS (5), PUMPKINS (0), SQUASH (5). Cucumber Beetle, Flea Beetle, Leafhopper, Pickleworm, Melonworm, Squash Bug; also Alternaria Blight, Angular Leaf Spot, Anthracnose, Blossom Blight, Downy Mildew, Gummy Stem Blight, Scab. Do not apply when foliage is wet creair or excessive humidity is expected during the next two days. Do not us on vatermelons in Florida.

ENDIVE (14), LETTUCE: Head (10), Leaf (14). Armyworm, Corn Earworm, Flea Beetle, Harlequin Bug. Imported Cabbageworm, Six-Spotted Leafhopper, Stink Bug, Tarnished Plant Bug; also Downy Mildew. Do not apply when foliage is wet or when excessive humidity or rain is expected during the next two days.

SPINACH (14), SWISS CHARD (14). Armywerm, Flee Beetle, Leafhopper, Stink

MAY 20 1914

UNDER ALL RUPENTICIDE ACT FOR ECONOMIC POISON REGILTER ED UNDER NO. 2/25-6 9 SUBJECT TO ATTACHED COMMENTS.



TONATO & VEGETABLE DUST



Bug, Tarn., hed Plant Bug; also Downy Mildew (Blue Mold), Leaf Spots, White Rust.

PEAS (0). Alfalfa Looper, Armyworm, Grasshopper, Leathopper; also Anthracnose (begin after blossoms open), Downy Mildew, Rust.

PEPPERS (0). CHmbing Cutworm, Colorado Potato Beetle, European Corn Borer, Fall Armywerm, Flea Beetle, Leafhopper. Stink Bug, Tomato Fruitworm, Tomato Hornworm, Ternished Plant Bug; 2130 Anthracnose, Bacterial Leaf Spot, Downy Mildew, Cercospora and other Leaf Spots, Phytophthora and other Fruit Rots.

Caution. Harmful if swallowed or inhaled. May cause irritation of nose, throat, skin. Do not breathe dust. In case of contact with skin or eyes wash thoroughly with water; for eyes get medical attention. Wash thoroughly after handling. Do not store near, or contaminate, feed or foodstuffs.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes

NOTICE. Buyer, or User, assumes all risks of use, storage or handling of this material not in strict accordance with the directions given on this label.

EPA Reg. No. 2125-69-AA Manufactured In U.S.A. by

SCIENCE PRODUCTS COMPANY Inc. Chicago, Illinois



as listed under Directions

TOMATOES and VEGETABLES

ACTIVE INGREDIENTS		
Carbaryl (1-naphthyl N-methylcarba	mate)	5.0%
Zineb (Zinc ethylene bisdithiocarba (metallic zinc equivalent - 1.65%	mate)	.7 .0°⁄3
INERT INGREDIENTS		88 .0°°
	Total	100.0%
Net Weight: 10	OZ.	
	3.5 gms.)	

*SEVIN - registered trademark of Union Carbide Corporation for carbaryl insecticide (1-naphthyl N-methylcarbamate)
U.S. Pat. Nos. 2,903,478, 3,009,855

CAUTION: KEEP OUT OF REACH OF CHILDREN See other cautions on back panel.