

Established 1884
HOFFMAN'S
GARDEN INSECTICIDE

This product is a concentrated, efficient formulation for use on tomatoes, cucumbers, melons, potatoes, beans, cole crops, sweet corn and peas; flowers, cabbage, broccoli, and kale; ornamental evergreens, shrubs, vines and apples, plums, blueberries, strawberries and raspberries.

ACTIVE INGREDIENTS	
Rotenone	1.11%
Other Cube Extractives	1.65%
Malathion—O, O-Dimethyl dithiophosphate of diethyl mercaptosuccinate ..	42.93%
Methylated Naphthalenes	32.41%
INERT INGREDIENTS	21.90%

DIRECTIONS 100.0%

Shake well before using.

For aiding in control of Mexican bean beetle on beans; flea beetle, potato beetle and leafhopper on potatoes; flea beetles on cabbage, broccoli and kale; thrips, leafhoppers and various aphids on cherries, plums, apples (may injure foliage on some varieties), beans, peas, tomatoes and strawberries; whitefly on melons; cucumber beetle on cucumbers; Japanese beetles on blueberries; mealy bug on raspberries; aphids, imported cabbage worm and cabbage looper on cabbage, broccoli and kale; aphids, leafhoppers and mealy bugs on flowers and aphids, bagworms and tent caterpillars on ornamentals and evergreens add 1 tablespoonful of this product to a gallon of water and cover both the upper and lower leaf surfaces, using 30 gallons per acre.

WARNING: Keep out of reach of children.

See other precautions on the side panel.

ACCEPTED
 2136-17
 SEP 4 1973
 UNDER THE FEDERAL INSECTICIDE ACT
 FUNGICIDE AND RODENTICIDE REGISTRATION
 FOR THE CONTROL OF CORN EARWORM LARVAE
 REGISTERED UNDER NO. 2136-17
 SUBJECT TO APPLICABLE COMMENTS.

For aiding in control of corn earworm larvae on corn, apply to silks as soon as they appear. Mix 1 tablespoonful of this product in 1½ gallons of water. Make 3 to 4 applications at 2 to 3 day intervals. Re-apply immediately after a rain.

For aiding in controlling cutworms in corn and tomato fields, add 2 tablespoonfuls of this product to 1 gallon of water and spray thoroughly the soil around the plants to be protected preferably in the late afternoon.

NOTE: Do not apply to beans after pods begin to form and do not feed the treated vines to livestock. Do not apply less than 7 days before harvest to other crops listed.

Do not reuse empty container. Wash thoroughly with soap and detergent and discard in a safe place.

WARNING: Do not get in the eyes and avoid breathing the mist. Poisonous if swallowed or absorbed thru the skin. Wash with soap and water after handling and before eating or smoking. In case of contact, wash immediately with soap and water. Wash all contaminated clothing before re-use.

CAUTION: This product is highly toxic to bees exposed to direct treatment or to a residue on crops. Do not apply or allow to drift to areas occupied by humans or beneficial animals. This product is poisonous to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

EPA Reg. No. 2136-17
 Net Fluid Contents 1 U. S. Quart
 Formulated by
J. L. Hoffman Co., Inc. Allentown, Pa. 18102