HOFFMAN'S
MALATHION 4 DUST

ACTIVE INGREDIENT:
Malathion (O-O-Dimethyl Dithiophosphate of Diethyl Mercaptosuccinate) 4%
INERT INGREDIENTS 96%
——
100%

As an aid in controlling horn flies on dairy cattle, after milking, dust 4 tablespoonfuls of this powder over the cows backs, necks and shoulders. This treatment will give good results for a week or more. If not washed off. Repeat application every two weeks or. as. necessary.

CAUTION: — Do not use the dust later than 5 hours before milking.

As an aid in controlling horn flies, on beef cattle and fleas and ticks on dogs and cats, apply thoroughly to animals and apply 1 to 2 lbs. per 1000 square feet of surface to pet quarters. Remove manure or debris before dusting. Repeat treatment within 3 to 4 weeks, if necessary.

Do not treat kittens or puppies under 4 weeks of age.

Do not treat animals under one month of age.

As an aid in controlling chicken red mites and northern fowl mites in the litter, work into the litter one pound of this product to every 50 square feet of floor area and mix the dust in the nest material. Individual birds may be dusted and be cleaned of all types of lice and mites.

As a dustbath for poultry, as an aid in controlling northern fowl mite and body lice, spread a pound of this product over the bottom of an 18 inch by 12 inch by 3 inch box and set where 30 hens can use it. Refill when the dust is used up.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Avoid inhaling the dust. Avoid contamination of food and feed stuffs. Avoid skin contact. Wash after handling. Avoid getting the dust into feed hoppers, drinking cups or water fountains. Do not reuse empty container. Destroy when empty.

CAUTION: This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning equipment, or disposal of wastes. Apply this product only as specified on this label.

NET WEIGHT   5 POUNDS

ACCEPTED WITH COMMENTS