### GENERAL INFORMATION

DRINOX H-34B, when used as directed, will help protect seeds and seedlings of wheat, barley, oats, corn, sorghum and cotton, from wireworms and seed-corn maggots. DRINOX H-34B, when used as directed, will help protect corn against attack from seed-corn beetles and Southern corn rootworms. It will also help to protect barley, oats and wheat from attack by false wireworms, and sorghum from attack by Kafir ants.

#### CAUTION

This product is toxic to fish and wildlife. Keep out of any body of water. Treated seeds exposed on soil surface will be hazardous to birds and other wildlife. Do not re-use empty drum. Destroy by perforating or crushing and burying in non-crop lands away from water supplies.

#### **PRECAUTIONS**

KEEP AWAY FROM HEAT AND OPEN FIRE. Flash point above 90 F. The crystals that may form at extremely low temperatures should disappear after the temperature of the liquid is increased and the container well shaken.

Treated seeds should be labeled as follows:


PO'SON TREATED—THIS SEED TREATED WITH HEPTACHLOR.

DO NOT USE FOR FOOD, FEED, OR OIL PURPOSES.

6225



# READY TO USE LIQUID SEED INSECTICIDE



CAUTION: Combustible—Keep away from heat and open flame.

May be fatal if swallowed. Do not get in eyes, on skin, or on clothing. Do not breathe vapor or fumes. Do not use for focd, feed or oil purposes.

#### ANTIDOTE

INTERNAL: Emetic of 1 tablespoonful mustard in a tunibler of water.

CALL A PHYSICIAN

EXTERNAL: In case of spillage on skin wash promptly with scap and water.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONS

**NET CONTENTS:** 



# NOR-AM.

AGRICULTURAL PRODUCTS, INC.

#### DIRECTIONS FOR USE

DRINOX H-348 is ready to use. It cannot be diluted with water or products containing water. It should be used in treaters which can apply the correct dosages shown below.

	DOSAGE DIRECTIONS Insects	Dosage fl. ozs./bushel
Seed		
Wheat	Wireworms, seed-corn maggots,	21/4
Barley	Wireworms, seed-corn maggots,	1 7/8
Oats	Wireworms, seed-corn maggots, alse wireworms	11/4
Corn	naggots, wireworms	11/4
Corn	Southern corn rootworms	21/4
Sorghum	Wireworms, seed-corn maggots, talse wireworms, Kafir ants	1%
Cotton	False wireworms, seed-corn inaggots, wireworms	4 fl. ozs./100 lbs.

Seeds should be cleaned and must be well cured before treatment.

#### IMPORTANT READ BEFORE PURCHASE OR USE

Purchase or use of this product shall constitute acceptance of the following terms of sale NOR-AM Agricultural Products, Inc. (Nt R AM) warrants that this product conforms to the chemical descriptions on this label and is reasonably it for the purposes specified hereom. By way of limitation but not in derogation of the directions and statements printed on this label, Selter makes no warranty and no representation and HOR AM makes no further warranty or representation expressed or implied, including but not limited to any warranty as to MERCHANTABILITY, and no agent of NOR AM is authorized to do so except in writing, with a specific reference to the foregoing warranty. Since neither Seller nor NOR AM has any control over the handling storage, use or conditions of use of this product, Buyer or user assumes all risks and liability arising from handling, storage or use of this product under abnormal conditions or not in strict compliance with the directions and precautionary statements hereon. All claims related to the purchase, handling, storage or use of this product shall be barred unless written notice is given to NOR-AM within 30 days after the loss is discovered. I liability of the Seller or NOR AM for any damages arising from a breach of this warranty shall be limited to exclude consequential commercial damages.

## DO NOT RE-USE CONTAINER

PRODUCT OF U.S.A.

PRINTED IN U.S.A.

