

**IMPORTANT:  
READ BEFORE PURCHASE OR USE**

Purchase or use of this product shall constitute acceptance of the following terms of sale. NOR-AM Agricultural Products, Inc. (NOR-AM) warrants that this product conforms to the chemical descriptions on this label and is reasonably fit for the purposes specified hereon. By way of limitation but not in derogation of the directions and statements printed on this label, Seller makes no warranty and no representation and NOR-AM makes no further warranty or representation, expressed or implied, including but not limited to any warranty as to MERCHANTABILITY, and no agent of NOR-AM is authorized to do so except in writing, with a specific reference to the foregoing warranty. Since neither Seller nor NOR-AM has any control over the handling, storage, use or conditions of use of this product, Buyer or user assumes all risks and liability arising from handling, storage or use of this product under abnormal conditions or not in strict compliance with the directions and precautionary statements hereon. All claims related to the purchase, handling, storage or use of this product shall be barred unless written notice is given to NOR-AM within 30 days after the loss is discovered. Liability of the Seller or NOR-AM for any damages arising from a breach of this warranty shall be limited to exclude consequential commercial damages.



E.P.A. REG. NO. 2139-99

**ACTIVE INGREDIENT:**

Formetanate hydrochloride\*.....92%

**INERT INGREDIENTS:**

8%

\*m-[[[(Dimethylamino)methylene]amino]phenyl methylcarbamate monohydrochloride



**POISON**



**DANGER**

**KEEP OUT OF REACH OF CHILDREN**

Not for use or storage in or around the home.

Call a physician immediately in case of exposure.

This product is poisonous if swallowed. Do not get in eyes, on skin or on clothing. Do not breathe dust or spray mist. Do not contaminate food and feedstuffs. Wear protective clothing, goggles and rubber (natural or synthetic) gloves when handling this material. Wash before eating or smoking. Wash contaminated clothing before re-use.

If exposure to dust or spray mist cannot be avoided wear a self-contained breathing apparatus.

**ANTIDOTES:**

If swallowed—give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Have patient lie down and keep quiet. Call a physician immediately.

If on skin—wash immediately with soap and water.

If in eyes—flush with water for at least 15 minutes and get medical attention.

**NOTICE TO PHYSICIAN:**

CARZOL SP is a carbamate which acts as a cholinesterase inhibitor.

Specific antidote—Atropine

Do not use 2-PAM.

SEE REAR LABEL FOR ADDITIONAL PRECAUTIONS AND DIRECTIONS FOR USE.

U.S. PAT. NO. 3336186  
3542853

NET CONTENTS: POUNDS

MANUFACTURED FOR

**NOR-AM®**

AGRICULTURAL PRODUCTS, INC.

20 N. WACKER DRIVE, CHICAGO, ILLINOIS 60606



2139-99

9-21-1972

1/5-


### DIRECTIONS FOR USE

CARZOL SP is completely soluble in water. Preparation of a premix or slurry before adding to the spray tank is not necessary. Add the desired amount of CARZOL SP directly to the spray tank 1/2 full of water with the agitator running. Then add the remainder of the water.

NOTICE: CARZOL SP is not stable in alkaline water. Do not prepare the spray mixture until just before application. Do not prepare more spray mixture than will be used within 4 hours.

CROP	PEST	APPLICATION
Oranges, lemons & tangerines	THrips in California & Arizona, only	Use 8 oz. CARZOL SP per 100 gallons of water. Apply for outside coverage in 100 to 300 gallons of water per acre (1/2 to 1 1/2 lbs. CARZOL SP per acre) depending on the size of the trees. Repeat application as needed (see note below).
	Red mite in California & Arizona, only	Use 8 oz. CARZOL SP per 100 gallons of water. Apply in a full coverage spray in 700 to 1000 gallons of water per acre (3 1/2 to 5 lbs. CARZOL SP per acre) depending on the size of the trees (see note below).
Oranges, lemons tangerines and grapefruit	Rust mite in Florida	Use 1 to 2 oz. CARZOL SP per 100 gallons of water. Apply in full coverage spray. Use 7 to 20 oz. CARZOL SP per acre depending on the size of the trees. Repeat application as needed.
	Rust mite in Texas	Use 2 to 4 oz. CARZOL SP per 100 gallons of water. Apply in full coverage spray. Use 10 to 20 oz. CARZOL SP per acre depending on the size of the trees. Repeat application as needed.
Apples and pears	European red mite, Two spotted mite	Use 4 to 8 oz. CARZOL SP per 100 gallons of water. Apply in full coverage sprays in 400 to 800 gallons of water per acre (1 to 4 lbs. of CARZOL SP per acre) depending on the size of the trees and severity of the mite infestation. Repeat application as needed (see note below)

3/5



225744  
04-043

$\frac{4}{5}$ 

Licensed by and Trademark\* of Schering A. G. Berlin, Bergkamen 

5/5

CPA-PA  
+ 0 0 0 0

ACCEPTED  
SEP 21 1972  
OF FUNGI  
FOR  
ED UNDER  
TO ATTACHED COMMENTS

2139-99