

2139-99

02/16/1973

1/2

Best Available Copy



Carzol[®] SP

MITICIDE / INSECTICIDE

E.P.A. REG. NO. 2139-99

ACTIVE INGREDIENT:

Formetanate hydrochloride* 92%

INERT INGREDIENTS: 8%

*m [((Dimethylamino) methylene) amino] phenyl methyl carbamate monohydrochloride



POISON DANGER



KEEP OUT OF REACH OF CHILDREN

Not for use or storage in or around the home.
Call a physician immediately in case of exposure.

This product is poisonous if swallowed. Do not get in eyes, on skin or on clothing. Do not breathe dust or spray mist. Do not contaminate food and feedstuffs. Wear protective clothing, goggles and rubber (natural or synthetic) gloves when handling this material. Wash before eating or smoking. Wash contaminated clothing before re-use.
If exposure to dust or spray mist cannot be avoided wear a self-contained breathing apparatus.

ANTIDOTES:

If swallowed—give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Have patient lie down and keep quiet. Call a physician immediately. If on skin—wash immediately with soap and water. If in eyes—flush with water for at least 15 minutes and get medical attention.

NOTICE TO PHYSICIAN:

CARZOL SP is a carbamate which acts as a cholinesterase inhibitor.
Specific antidote—Atropine.
Do not use 2-PAM.

See Rear Label For Additional Precautions and Directions For Use.

U.S. PAT. NOS. 3336186—3542853

NET CONTENTS: 2½ POUNDS



MANUFACTURED FOR
NOR-AM.

AGRICULTURAL PRODUCTS, INC.
CHICAGO, ILLINOIS 60606

$\frac{2}{2}$

FUNCTIONAL AND RELEVANT ACT
FOR A ... SUBJECT
TO ATTACHED COMMENTS

CARZOL SP is completely soluble in water. No special pre-treatment is necessary before adding to the spray tank. Add the desired amount of CARZOL SP to the spray tank 1/2 full of water with the agitator running. Then add the remainder of the water.

CROP	PEST	APPLICATION
ORANGES, LEMONS AND TANGERINES	THRUST MITE IN CALIFORNIA & ARIZONA, ONLY	Use 8 oz. CARZOL SP per 100 gallons of water. Apply for mite coverage in 100 to 300 gallons of water per acre (½ to 1½ lbs. CARZOL SP per acre) depending on the size of the trees. Repeat application as needed (see note below).
	RED MITE IN CALIFORNIA & ARIZONA, ONLY	Use 8 oz. CARZOL SP per 100 gallons of water. Apply in a full coverage spray in 700 to 1000 gallons of water per acre (3¼ to 5 lbs. CARZOL SP per acre) depending on the size of the trees (see note below.)
ORANGES, LEMONS, TANGERINES AND GRAPEFRUIT	RUST MITE IN FLORIDA	Use 1 to 2 oz. CARZOL SP per 100 gallons of water, apply in full coverage spray. Use 7 to 20 oz. CARZOL SP per acre depending on the size of the trees. Repeat application as needed.
	RUST MITE IN TEXAS	Use 2 to 4 oz. CARZOL SP per 100 gallons of water, apply in full coverage spray. Use 10 to 20 oz. CARZOL SP per acre depending on the size of the trees. Repeat application as needed.
APPLES AND PEARS	EUROPEAN RED MITE, TWO SPOTTED MITE	Use 4 to 8 oz. CARZOL SP per 100 gallons of water. Apply in full coverage sprays in 400 to 800 gallons of water per acre (1 to 4 lbs. of CARZOL SP per acre) depending on the size of the trees and severity of the mite infestation. Repeat application as needed (see note below).

To avoid excessive residues, observe the following limitations: *CITRUS*: Do not apply more than 5 lbs. CARZOL SP per acre in a 12 month period. Allow 30 days between applications and 7 days before harvest. *Other Crops*: Do not apply more than 4 lbs. CARZOL SP per acre in one crop season. Do not apply less than 7 days before harvest.

This product is toxic to birds and other wildlife.

Do not contaminate any body of water with this material. Apply only as specified on this label. Dispose of left-over material or empty containers by burning or burying in non-crop land away from water supplies. Stay out of smoke. Do not reuse containers. Call 1-800-SP-4-666 for more information. Always follow instructions on the label. For more information, call 1-800-SP-4-666.



Another person who has been involved in the

NOR-AM.

AGRICULTURAL PRODUCTS, INC.
CHICAGO, ILLINOIS